

in a cardboard box equipped with a clear plastic emergence vial. The boxes were kept at 21°C and 70% RH. Parasites began to emerge on 18 July; they were removed from the vials twice a day and, to ensure mating, were kept in a screen cage (30 x 30 x 56 cm), with raisins and water provided until release. The parasites were released twice a week from 21 July to 15 August inclusive at 9.7 km outside of St. John's (52°47' long. and 47°37' lat.) in a sawfly-infested plantation comprising Scots pine, *Pinus sylvestris* L., and Jack pine, *Pinus banksiana* Lamb. in about equal proportion. Totals of 631 males and 376 females were released.

Parasitism by *P. basizonus* was determined on naturally occurring cocoons and on planted cocoons. Naturally occurring cocoons were collected from 20 litter and soil samples (30 x 30 x 8 cm) taken from under the crown of sawfly-infested trees. Three trays (20 x 20 x 3 cm, covered with 0.5 cm mesh screen on the bottom and top) were placed at the release site with laboratory-reared cocoons to expose them to attack by *P. basizonus*. Each tray contained 50 sawfly cocoons, which were replaced once a week with fresh ones. Cocoons totalling 600 were exposed to parasites in this manner. These and the cocoons obtained from the soil samples were reared in the laboratory at 21°C and 70% RH until October, then stored at 2°C for 3 mo, and then reared again.

Only five sawfly cocoons were obtained from the 20 soil samples, and no parasites emerged from them. Of the 600 sawfly cocoons exposed to parasites in the trays, 30% emerged as sawfly adults, 37% died of unknown causes in the cocoon stage, 11% were parasitized by *Mastrus aciculatus* (Prov.), and 22% were parasitized by *P. basizonus*.

M. aciculatus, an ichneumonid, has been reared from the European pine sawfly in Ontario (Griffiths, Can. Entomol. 91:501-512, 1959), but this is the first record of it from this host in Newfoundland. The recovery of *P. basizonus* from such a large percentage of the planted cocoons indicates that at least one generation of parasites has survived. However, the successful establishment of this parasite may be verified only through several years of monitoring.

Technical assistance in this study was provided by D.L. Oliver and K.E. Pardy.—Imre S. Otvos, Newfoundland Forest Research Centre, St. John's, Nfld., and K.J. Griffiths, Great Lakes Forest Research Centre, Sault Ste. Marie, Ont.

Biological Observations on Overwintering Larvae of the Large Aspen Tortrix in Alberta.—The large aspen tortrix, *Choristoneura conflictana* (Wlk.), is a serious pest of aspen in North America. The life history of this insect and the damage caused by it have been reported in Canada (Prentice, Can. Entomol. 87:461-473, 1955), in California (Wickman, J. Econ. Entomol. 56:593-596, 1963), and in Alaska (Beckwith, USDA Forest Serv., Pac. Northwest Forest Range Exp. Stn. Res. Note PNW-81, 1968). Eggs of the large aspen tortrix are laid from mid-June to early July and hatch in July. First-instar larvae feed on leaf surfaces. In late July or August they undergo their first molt and then hibernate; second-instar larvae apparently do not feed the first summer, remaining in hibernation until new growth starts in the spring. Pupation usually occurs in early June, but this depends on the climate. This note reports on the site and parasites of overwintering larvae of the large aspen tortrix in Alberta.

According to Prentice (1955), in Manitoba and Saskatchewan the second-instar larvae overwinter in bark crevices and moss at the base of living trees. In their current studies on the biological control of *C. conflictana* in Alberta, however, W.G.H. Ives and J.A. Muldrew (pers. commun.) have noted that overwintering larvae were found beneath the slightly loosened bark of the trunk and branches of dead trembling aspen. Their observations agree with those of Wickman (1963) in California and Beckwith (1968) in Alaska.

The parasites associated with the large aspen tortrix in Manitoba and Saskatchewan have been listed by Prentice (1955), who also produced survey rearing records for New Brunswick, Ontario, and Alberta. Three additional species of parasites not listed by Prentice (1955) have been identified in Alberta. They were recovered from overwintering second-instar larvae collected by Ives and Muldrew in early March 1978 at Hondo, Alta., and sent to J.C. Cunningham, Forest Pest Management Institute, Sault Ste. Marie, Ont., for virus host range studies. About 2,500-3,000 surplus larvae were reared on an artificial diet (Grisdale, Can. Entomol. 105:1553-1557, 1973) in 21.3 mL cream cups kept at 23.9°C with a relative humidity of 50-55% (Cunningham et al., Can. Entomol. 105:767-773, 1973). An approximately 50% level of parasitism was noted for these second-instar *C. conflictana* larvae from Alberta (D. Grisdale, pers. commun.). The

parasites were returned to the Edmonton laboratory, where they were identified as *Macrocentrus iridescens* French (Hymenoptera: Braconidae), *Glypta inversa* Cress. (Hymenoptera: Ichneumonidae), and *Agathis annulipes* (Cres.) (Hymenoptera: Braconidae). Collecting overwintering larvae in late winter and rearing them revealed also that the three parasites attacked the large aspen tortrix larvae in their first year.

The most common parasite was *M. iridescens*, followed by *G. inversa* and then by *A. annulipes*. The recovery of the latter two species in Alberta confirms observations by Torgersen and Beckwith (Can. Entomol. 106:1247-1265, 1974) that *G. inversa* and *A. annulipes* attack first- or second-instar larvae of *C. conflictana*, and the rearing results indicate further that *M. iridescens* also attacks first- or second-instar larvae of *C. conflictana*.

It is interesting to note that Torgersen and Beckwith (1974) found *G. inversa* to be the most common parasitoid attacking early larvae of the large aspen tortrix in interior Alaska but did not recover any *M. iridescens*, which was predominant in Alberta. The predominance of *Macrocentrus* over *Glypta* in Alberta is suggestive of the relationship between the same two parasitic genera of the oriental fruit, *Grapholitha molesta* (Busck), in Ontario. Steenburgh and Boyce (Annu. Rep. Entomol. Soc. Ont. 69:65-74, 1938) observed that as *Macrocentrus ancylivorus* Roh. became more plentiful, *Glypta ruficollaris* Cress. became increasingly less important. This may offer an explanation for the absence of *Macrocentrus* and the predominance of *Glypta* in the interior of Alaska.—H.R. Wong, Northern Forest Research Centre, Edmonton, Alta.

High Populations of a Carabid Beetle Associated with Spruce Budworm.—Numerous large, black carabid beetles (*Calosoma frigidum* Kby.) were noted in two white spruce (*Picea glauca* [Moench] Voss) plantations near Sault Ste. Marie, Ont., in early June 1977. The beetles were seen eating late-instar spruce budworm (*Choristoneura fumiferana* [Clem.]) larvae, and it is possible that they played some part in reducing budworm populations in the plantations.

The two plantations were visited at least once a week from mid-May to the end of July in connection with other studies. Plantation A (250 ha) was planted with white spruce and, in one location, with Norway spruce (*P. abies* [L.] Karst.), in 1925-28; plantation B (12 ha) was planted during the same period with white spruce only. Both plantations now contain considerable jack pine (*Pinus banksiana* Lamb.) and occasional white pine (*P. strobus* L.). The dominants are 15-17 m tall, but there are numerous small openings dominated by *Vaccinium* species. Located on sandy, gravelly river terraces on either side of the Goulais River, a few kilometers north of Searchmont, these are the only white spruce plantations in the vicinity. Natural stands dominated by balsam fir (*Abies balsamea* [L.] Mill.) and white spruce occur in small patches along river valleys nearby, but most of the area is covered by hardwood with a low proportion of white spruce.

Twenty-five 45 cm branch tips were taken from each plantation on 26 May, 1977, and examined for spruce budworm larvae. In plantation A counts averaged 32.0 larvae/branch tip and in B they averaged 31.8. At this time, fourth-, fifth-, and sixth-instar larvae accounted for 25%, 45%, and 30%, respectively, of the total. On 30 June, pupal populations averaged 0.08/45 cm branch tip in A and 0.04 in B—99.8 and 99.9% less, respectively. Egg populations, sampled on 27 July, averaged 0.86 egg masses/45 cm branch tip in A and 0.71 in B (equivalent to approximately 85 and 70/10 m², respectively).

The carabids were first noticed on 6 June, on the lower stems of the spruce. A quantitative assessment was made on 16 June. Initially, attempts were made to obtain estimates of density by beating foliage over drop sheets. However, only five beetles were dislodged from 50 trees. Several beetles remained on the foliage after beating, and more were observed on the higher foliated branches that were out of reach of the beaters than on the dead lower branches. A visual assessment was therefore made on both foliated and defoliated portions of trees. Fifty trees 8-10 m high, all in plantation B, were examined. Observations were restricted to the base and to branches up to 2 m above ground. The trees were categorized as (a) having no living needles on the observed branches, (b) having foliage on the periphery of some branches, (c) having well foliated branches down to the ground. Beetle counts are given in Table 1. Fifty-two beetles were seen on the 50 trees—six on the tree boles and 46 on branches. As many as six were seen on one tree, but 25 trees had no beetles, although 20 of the 35 well-foliated trees had at

bi-monthly research notes

Vegetative Propagation of Trembling Aspen with Auxins

Microsporogenesis in Larix Laricina in eastern Newfoundland

Conifer Seed Pathogenicity Tests with Forest Cup Fungi

Cocoon Parasite of the European Pine Sawfly Introduced into Newfoundland from Ontario

Biological Observations on Overwintering Larvae of the Large Aspen Tortrix in Alberta

High Populations of a Carabid Beetle Associated with Spruce Budworm

Pine Oil Delays Attack of Ambrosia Beetles on Piled Log Sections

Chemical Control Trials on the Box Elder Twig Borer in Alberta

Vol. 35, No. 4, JULY-AUGUST 1979



Environment
Canada

Environnement
Canada

Forestry
Service

Service
des forêts