

1979

THIS FILE COPY MUST BE RETURNED

TO: INFORMATION SECTION,  
NORTHERN FOREST RESEARCH CENTRE,  
5320-122 STREET,  
EDMONTON, ALBERTA,  
T6H 3S5



Interim Report on  
Physiology and Mechanisms of Air-Borne  
Pollutant Injury to Vegetation,  
1975 to 1978

Project VE 3.3  
AOSERP Report 45  
January 1979

Sponsored jointly by



Environment  
Canada

Environnement  
Canada

15th Floor, Oxbridge Place  
9820 - 106 Street  
Edmonton, Alberta, Canada  
T5K 2J6

ALBERTA OIL SANDS ENVIRONMENTAL RESEARCH PROGRAM

RESEARCH REPORTS

---

These research reports describe the results of investigations funded under the Alberta Oil Sands Environmental Research Program, which was established by agreement between the Governments of Alberta and Canada in February 1975 (amended September 1977). This 10-year program is designed to direct and co-ordinate research projects concerned with the environmental effects of development of the Athabasca Oil Sands in Alberta.

A list of research reports published to date is included at the end of this report.

Enquiries pertaining to the Canada-Alberta Agreement or other reports in the series should be directed to:

Alberta Oil Sands Environmental Research Program  
15th Floor, Oxbridge Place  
9820 - 106 Street  
Edmonton, Alberta  
T5K 2J6  
(403) 427-3943

---

Interim Report on Physiology and  
Mechanisms of Air-Borne Pollutant Injury  
to Vegetation, 1975 to 1978

Project VE 3.3

AOSERP Report 45

This report may be cited as:

Malhotra, S.S. 1979. Interim report on physiology and mechanisms of air-borne pollutant injury to vegetation, 1975 to 1978. Prep. for the Alberta Oil Sands Environmental Research Program by Canadian Forestry Service, Northern Forest Research Centre. AOSERP Report 45. 38 pp.

The Hon. D.J. Russell  
Minister of the Environment  
222 Legislative Building  
Edmonton, Alberta

and

The Hon. L. Marchand  
Minister of State for the Environment  
Fisheries and Environment Canada  
Ottawa, Ontario

Sirs:


Enclosed is the report "Interim Report on Physiology and Mechanisms of Air-Borne Pollutant Injury to Vegetation, 1975 to 1978".

This report was prepared for the Alberta Oil Sands Environmental Research Program, through its Vegetation Technical Research Committee (now part of the Land System), under the Canada-Alberta Agreement of February 1975 (amended September 1977).

Respectfully,



W. Solodzuk, P. Eng.  
Chairman, Steering Committee, AOSERP  
Deputy Minister, Alberta Environment



A.H. Macpherson, Ph.D.  
Member, Steering Committee, AOSERP  
Regional Director-General  
Environmental Management Service  
Fisheries and Environment Canada

INTERIM REPORT ON PHYSIOLOGY AND  
MECHANISMS OF AIR-BORNE POLLUTANT INJURY  
TO VEGETATION, 1975 to 1978

DESCRIPTIVE SUMMARY

BACKGROUND AND PERSPECTIVE

Full understanding of plant injury thresholds requires a knowledge of the extent and types of injury to be expected at different exposures and conditions as injury development is a continuum, not a "present" or "not present" condition.

The main goal of this research project is the development of a field test for the detection of previsual symptoms of air-borne pollutant damage. It is necessary to study the physiology of plant response to effluents (including cellular, structural, and bio-chemical changes) since this is the level at which effects can first be detected. Since plants often experience sub-lethal exposures which appear to result in long term debilitation of physiological activity, the detection of previsual symptoms is necessary to allow economic and ecological evaluation of the real effects of chronic low level exposure to effluents.

Since development of visual symptoms is often indicative of debilitation so extensive that recovery is doubtful, long term biomonitoring of the ecosystem should be based on previsual symptoms which occur before recovery is doubtful.

This research project has been designed and carried out in conjunction with project VE 3.1 (Symptomology and Threshold Levels of Air Pollutant Injury to Vegetation, 1975 to 1978) and project VE 3.4 (Ecological Benchmarking and Biomonitoring for Detection of Air-Borne Pollutant Effects on Vegetation and Soils, 1975 to 1978).

ASSESSMENT

The report "Interim Report on Physiology and Mechanisms of Air-Borne Pollutant Injury to Vegetation, 1975 to 1978", prepared by S.S. Malhotra (Northern Forest Research Centre, Canadian Forestry Service) has been reviewed and accepted by the Alberta Oil Sands Environmental Research Program.

In view of the value of the document, Program Management of AOSERP recommended that the report be published and made available to other AOSERP researchers and the public as soon as possible.


The report is comprehensive and discusses results obtained under controlled conditions and results from material subjected to field impingement. Preliminary conclusions are presented.

The content of this report does not necessarily reflect the views of Alberta Environment, Fisheries and Environment Canada, or the Alberta Oil Sands Environmental Research Program. The mention of trade names for commercial products does not constitute an endorsement or recommendation for use.



---

S.B. Smith, Ph.D.  
Program Director  
Alberta Oil Sands Environmental  
Research Program



---

R.A. Hursey, Ph.D.  
Research Manager  
Land System

INTERIM REPORT ON PHYSIOLOGY AND  
MECHANISMS OF AIR-BORNE POLLUTANT INJURY  
TO VEGETATION, 1975 to 1978

by

S.S. MALHOTRA  
Northern Forest Research Centre  
Canadian Forestry Service

for

ALBERTA OIL SANDS ENVIRONMENTAL  
RESEARCH PROGRAM

Project VE 3.3

January 1979

TABLE OF CONTENTS

	Page
DECLARATION . . . . .	ii
LETTER OF TRANSMITTAL . . . . .	iii
DESCRIPTIVE SUMMARY . . . . .	iv
LIST OF TABLES . . . . .	xi
LIST OF FIGURES . . . . .	xii
ABSTRACT . . . . .	xiii
ACKNOWLEDGEMENTS . . . . .	xiv
1. INTRODUCTION . . . . .	1
2. MATERIALS AND METHODS . . . . .	4
2.1 Growth Conditions . . . . .	4
2.2 SO <sub>2</sub> Treatment . . . . .	4
2.2.1 Aqueous SO <sub>2</sub> Treatment . . . . .	6
2.2.2 Gaseous SO <sub>2</sub> Treatment . . . . .	6
2.3 Pigment Related Assays . . . . .	7
2.3.1 Magnesium Determination . . . . .	7
2.3.2 Pigment Extraction . . . . .	7
2.3.3 Chlorophyllase Preparation . . . . .	8
2.4 Electron Microscopy . . . . .	8
2.5 Glycolipid Determination . . . . .	8
2.6 Soluble Sugar Determination . . . . .	8
2.7 Tissue Homogenate Preparation . . . . .	9
2.8 Enzyme Assays . . . . .	9
2.8.1 Acid Phosphatase . . . . .	9
2.8.2 Malate Dehydrogenase . . . . .	9
2.9 Lipid Biosynthesis . . . . .	10
3. RESULTS AND DISCUSSION . . . . .	11
3.1 Effects of Aqueous SO <sub>2</sub> on Total Chlorophyll Content . . . . .	11
3.1.1 Chlorophyll Breakdown . . . . .	11
3.1.2 Effect on Various Pigments . . . . .	13
3.1.3 Effect on Chlorophyllase Activity . . . . .	17
3.2 Ultrastructural Studies . . . . .	19
3.3 Effect of SO <sub>2</sub> on Tissue Glycolipids and Permeability Properties . . . . .	20
3.4 Effect of Gaseous SO <sub>2</sub> on Phosphatase Activity . . . . .	24

TABLE OF CONTENTS (CONCLUDED)

	Page
3.5	Effect of Gaseous SO <sub>2</sub> on Malate Dehydrogenase . . . . 26
3.6	Effect of Gaseous SO <sub>2</sub> on Lipid Biosynthesis . . . . . 28
3.6.1	Recovery of Fumigated Seedlings . . . . . 28
3.7	Biochemical Analyses on Field Samples . . . . . 30
4.	CONCLUSIONS . . . . . 32
5.	REFERENCES CITED . . . . . 33
6.	AOSERP RESEARCH REPORTS . . . . . 36



LIST OF TABLES

	Page
1. The Effect of Various Concentrations of SO <sub>2</sub> on Total Chlorophyll Contents of Pine Needles . . . . .	12
2. The Effect of Various Concentrations of Aqueous SO <sub>2</sub> on Mg <sup>++</sup> Loss from Pine Needle Chlorophyll . . . . .	15
3. The Effects of Gaseous SO <sub>2</sub> on Acid Phosphatase Activity in Young and Old Needles of Jack Pine Seedlings and its Ability to Recover in a SO <sub>2</sub> -free Atmosphere . . . . .	25
4. The Effects of SO <sub>2</sub> on Malate Dehydrogenase Activity from Pine Needles . . . . .	27
5. The Effects of Gaseous SO <sub>2</sub> on Lipid Biosynthesis in Developing Needles from Jack Pine Seedlings and their Ability to Recover in a SO <sub>2</sub> -free Environment . . . . .	29

LIST OF FIGURES

	Page
1. The AOSERP Study Area . . . . .	3
2. Map of the Area Around the Great Canadian Oil Sands Plant Showing Sites A, B, and C where Vegetation Samples for Biochemical Analyses were Collected . . . . .	5
3. Different Mechanisms of Chlorophyll Breakdown . . . . .	14
4. The Effect of Aqueous SO <sub>2</sub> on Chlorophyll <u>a</u> (A), Chlorophyll <u>b</u> (B), Chlorophyllide <u>a</u> (C), Chlorophyllide <u>b</u> (D), Phaeophytin <u>a</u> (E), and Phaeophytin <u>b</u> (F) . . . . .	16
5. The Effect of Aqueous SO <sub>2</sub> on the Activity of Chloro- phyllase . . . . .	18
6. The Effect of Aqueous SO <sub>2</sub> on the Levels of Glycolipids in Fully Developed and Young Pine Needles . . . . .	21
7. The Effect of Aqueous SO <sub>2</sub> on the Release of Soluble Sugars from Fully Developed and Young Pine Needles . . . . .	23

ABSTRACT

Biochemical studies conducted under controlled conditions suggested that low concentrations of  $\text{SO}_2$  which normally do not produce any visual symptoms on the foliage cause injury to forest vegetation by altering pigment (chlorophyll, phaeophytin and chlorophyllide) metabolism, by inhibiting lipid synthesis and activities of various important enzyme systems and by causing ultrastructural disorganization of cellular membranes.

In general, aqueous and gaseous  $\text{SO}_2$  produced results very similar to each other. Plants, when fumigated with  $\text{SO}_2$  at the ambient air quality standards, exhibited various biochemical responses that can have a deleterious effect on the normal growth and yield of vegetation. However, 24-48 h after transferring the fumigated plants to an  $\text{SO}_2$ -free environment, there was a considerable recovery of such functions.

The results obtained so far clearly indicated that chlorophyll/phaeophytin ratio,  $\text{Mg}^{++}$  content, lipid biosynthesis and acid phosphatase and malate dehydrogenase activities of plant foliage are very sensitive indicators of  $\text{SO}_2$  concentrations. Using such techniques, an attempt was made to detect pre-visual  $\text{SO}_2$  injury to jack pine and white spruce from the field impingement area (immediate vicinity of GCOS plant). Although a gradient in sulphur content of the collected material was evident, there was no appreciable difference in any of the biochemical and physiological functions that responded to  $\text{SO}_2$  under laboratory conditions. However, more experimentation needs to be conducted with some of the sensitive species of vegetation to determine air pollution effects on forest vegetation.

ACKNOWLEDGEMENTS

This research project VE 3.3 was funded by the Alberta Oil Sands Environmental Research Program, a joint Alberta-Canada research program established to fund, direct, and co-ordinate environmental research in the Athabasca Oil Sands area of north-eastern Alberta. The author thanks E. Hargesheimer, J. Shuya, and P. Hurdle for technical assistance.

## 1. INTRODUCTION

The growth and yield of vegetation are the direct results of physiological and biochemical activities within the tissues. Environmental factors such as temperature, moisture, light, etc. play an important role in determining the rate of metabolic activity. It has been suggested that air pollutants can affect tree metabolism either temporarily or permanently depending upon concentration and length of exposure (Brandt and Heck 1968; Saunders and Wood 1973; Ziegler 1975; Malhotra and Hocking 1976).

Sulphur dioxide ( $\text{SO}_2$ ) is one of the major air pollutants released from the industrial activity in the Athabasca Oil Sands area and may have the potential to cause damage to the forest vegetation and soils. Other air pollutants released along with  $\text{SO}_2$  may also influence such damage. There seems to be a tendency to relate phytotoxicity of pollutants only to visual symptom development (Malhotra and Hocking 1976). However, in many greenhouse studies,  $\text{SO}_2$  has been shown to reduce the growth and yield of vegetation in the absence of visual symptoms (Bleasdale 1952; Tingey *et al.* 1971). This may be the result of disturbances at the ultrastructural or biochemical levels or both.

There are a number of physiological and biochemical events in plant metabolism that are interfered with before the development of visual symptoms of  $\text{SO}_2$  phytotoxicity. Once injury to vegetation starts to occur at the visual symptom level, it may be irreversible. It is, therefore, essential that the early detection of air pollutant injury be carried out at biochemical and ultrastructural levels.

The only information available in the literature to determine previsual  $\text{SO}_2$  injury to forest species and to establish threshold levels of  $\text{SO}_2$  injury originates from other countries. However, due to considerable differences in vegetation type, climate and species present, such information is not directly applicable to the North American boreal forest species.

The objectives of this study are: (1) to study biochemical or physiological threshold levels of air pollutant injury to vegetation; (2) to develop sensitive and reliable methods for detecting air pollutant effects to forest vegetation prior to visual symptom development; and (3) to develop methods for selecting revegetation species on the basis of their true tolerance to air pollutants. In order to estimate the phytotoxic effects of various pollutants it is essential that the above objectives be tested first with each pollutant individually and then in combinations. The results described in this report relate mainly to the effects of SO<sub>2</sub> phytotoxicity; however, studies on combined effects of SO<sub>2</sub> with other pollutants are currently underway.

In the earlier phase of this program when the facilities for fumigating plant material with gaseous SO<sub>2</sub> were not available, research was carried out with aqueous SO<sub>2</sub> to understand the nature of phytotoxicity at the ultrastructural and biochemical levels. The results to date clearly suggest that the aqueous and gaseous SO<sub>2</sub> produce very similar effects.

Several techniques were developed and utilized to detect air pollution injury to forest species around the Alberta Oil Sands Environmental Research Program (AOSERP) study area (Figure 1). Further research is being conducted to establish threshold levels of air pollution injury to forest species. Because of the interim nature of this report, the statements made herein are subject to revision upon completion of the project and release of the final report.




Figure 1. The AOSERP study area.

## 2. MATERIALS AND METHODS

Most of the research in this project was conducted in the laboratory, greenhouse and fumigation chamber facilities located at the Northern Forest Research Centre, Edmonton. Field specimens to assess the impact of air pollution to jack pine and white spruce were collected from the impingement area at distances of 2.8 (A), 4.2 (B) and 8.3 (C) km south of the Great Canadian Oil Sands (GCOS) plant as shown in Figure 2. These sites were within a sulphur deposition gradient determined from a snow analysis survey (Barrie and Whelpdale 1978).

### 2.1 GROWTH CONDITIONS

Jack pine seeds (*Pinus banksiana* Lamb.) were treated with 3% H<sub>2</sub>O<sub>2</sub> for 10 min after which they were washed with distilled water and dried on paper towels. The seeds were planted in cavity trays in peat moss and irrigated. The trays were placed in the greenhouse and the seedlings grown under supplemented natural lighting (photoperiod 16 h). The seedlings were fertilized once a week with nutrient solution and watered as required.

### 2.2 SO<sub>2</sub> TREATMENT

It has been suggested that after penetration into plant tissues, gaseous SO<sub>2</sub> becomes hydrated, and it is the hydrated species that affects the various cellular processes (Hocking and Hocking 1977). Therefore, in our earlier studies, when the fumigation facilities were unavailable, we studied the effects of aqueous SO<sub>2</sub> on pigment metabolism and lipid biosynthesis. After acquiring the fumigation facilities in 1977, however, we investigated the effect of gaseous SO<sub>2</sub> on the activities of various important enzymes and lipid biosynthesis.

In equating the effects of gaseous and aqueous SO<sub>2</sub> on the biological activity of the spores of the fungus *Diplocarpon rosae*, Saunders (1966) observed that a X1000 higher concentration of SO<sub>2</sub> had to be employed in the aqueous phase than in the






Figure 2. Map of the area around the Great Canadian Oil Sands plant showing sites A, B and C where vegetation samples for biochemical analysis were collected.

gaseous phase. The above relationship is, however, dependent upon temperature and  $\text{SO}_2$  concentrations. Recently, Hocking and Hocking (1977) inferred that under moist conditions the dissolved unreacted species of  $\text{SO}_2$  ( $\text{SO}_2 \cdot 6\text{H}_2\text{O}$ ) would be the most likely biologically active species for plant injury. The concentration of dissolved unreacted  $\text{SO}_2$  at  $25^\circ\text{C}$  for 1, 10 and 100 ppm total  $\text{SO}_2$  in water was calculated to be 0.0012, 0.011 and 0.092 ppm, respectively or about 1/1000 of the total  $\text{SO}_2$  in water.

### 2.2.1 Aqueous $\text{SO}_2$ Treatment

Aqueous  $\text{SO}_2$  stock solution (1000 ppm) was prepared by bubbling anhydrous  $\text{SO}_2$  through a known quantity of 0.1 M Tris buffer (pH 7.2) until a desired gain of weight was reached (Puckett *et al.* 1973). Various concentrations (10, 25, 50, 100, 250, 500 and 1000 ppm) of aqueous  $\text{SO}_2$  were prepared by diluting the stock solution. An HCl control adjusted to the same pH as that obtained by 500 ppm  $\text{SO}_2$  (3.95–24 h after preparation of aqueous  $\text{SO}_2$  solution) was also prepared. Representative samples from excised pine needles from a population of 15–20 seedlings were incubated in these solutions under  $200 \mu\text{E m}^{-2}\text{s}^{-1}$  light source for varying lengths of time at  $23^\circ\text{C}$  and then utilized for experimental purposes.

### 2.2.2 Gaseous $\text{SO}_2$ Treatment

Fumigations were carried out in small acrylic cuvettes using a continuous flow-through system. Flow rates for air and  $\text{SO}_2$  were metered and controlled using rotameters and valves before being mixed and introduced into the cuvettes. Further mixing within the cuvettes was achieved by internally mounted fans near the injection ports. The concentrations of  $\text{SO}_2$  at the input and output were monitored by means of a Philips PW9700  $\text{SO}_2$  analyzer.

Pine seedlings (15–20 seedlings) were gently pulled out of the styrofoam trays and inserted into vertically slitted

neoprene stoppers so that the foliage extended through the top of the stopper and the peat plugs containing the roots extended through the bottom. The stoppers holding seedlings were inserted through holes cut in the bottom of the cuvette so that the slit closed firmly around the stem and sealed the hole. The cuvette was placed over a water tank filled enough to keep the peat plugs moist for the duration of the experiment. The access ports of the cuvettes were sealed to avoid any leakage. The air flow was  $10\text{ l min}^{-1}$  and  $\text{SO}_2$  flow was adjusted to give the desired concentration at the output of the cuvette.

The cuvettes were placed inside a controlled environmental chamber that maintained uniform light and temperature conditions. Fumigations were carried out at an isothermal temperature of  $22^\circ\text{C}$  with light intensity at approx.  $280\text{ }\mu\text{E m}^{-2}\text{s}^{-1}$  (16 h photoperiod). Relative humidity measured in the output gas stream varied between 60 and 80%.

## 2.3 PIGMENT RELATED ASSAYS

### 2.3.1 Magnesium Determination

Loss of magnesium from excised needles (5 g) treated with aqueous  $\text{SO}_2$  (22 h) was determined by means of Perkin Elmer atomic absorption spectrophotometer.

### 2.3.2 Pigment Extraction

Needles (5 g) treated with aqueous  $\text{SO}_2$  (22 h) were gently ground by pestle and mortar with the gradual addition of 20 ml of cold 100% acetone. The homogenate was filtered and the residue was washed twice with 80% acetone. The contents of various pigments were quantitatively determined spectrophotometrically according to the method of White *et al.* (1963).

### 2.3.3 Chlorophyllase Preparation

Aqueous SO<sub>2</sub>-treated needles (5 g) for 22 h were thoroughly homogenized. The homogenate was filtered and residue was washed as above. The residue thus obtained was further washed twice with 20 ml of absolute ethanol-anhydrous ether (3:1, V/V) and twice with 20 ml of anhydrous ether. The residue was dried at room temperature overnight and used for chlorophyllase assay (Holden 1961).

### 2.4 ELECTRON MICROSCOPY

Aqueous SO<sub>2</sub> treated (22 h) pine needle segments were fixed overnight at 4°C in 3% glutaraldehyde made up in 0.1 M phosphate buffer (pH 7.0). After fixation, the material was washed twice with the buffer and then post-fixed for 2-4 h in 2% osmium tetroxide dissolved in the phosphate buffer. After rinsing the material with the phosphate buffer it was dehydrated through a graded series of ethanol (50, 70, 85, 95, 100%). The excised needles were then infiltrated (under vacuum) with propylene oxide-araldite mixture (1:1, V/V) for 36 h. After infiltration, the sections were embedded in araldite and polymerised. The sections were cut, stained in 0.27% lead citrate for 2-3 min and examined under an electron microscope.

### 2.5 GLYCOLIPID DETERMINATION

Total lipids were extracted from the aqueous SO<sub>2</sub> treated (22 h) tissues and evaporated to dryness (Folch *et al.* 1957). These lipids were dissolved in chloroform and an aliquot was analyzed for the quantitative analysis of galactolipids and sulpholipids by thin layer chromatography on silica gel (Roughan and Batt 1968).

### 2.6 SOLUBLE SUGAR DETERMINATION

After the incubation (22 h) of excised pine needles in aqueous SO<sub>2</sub> the solution was removed and the tissues were washed

2-3 times with distilled water. The incubation solution and the washings from each sample were combined for soluble sugar analysis. HCl was added to an aliquot of the combined solution (final HCl conc. 3%) and centrifuged to remove any residue. The soluble sugars in the supernatant layer were determined according to the method of Dubois *et al.* (1956).

## 2.7 TISSUE HOMOGENATE PREPARATION

Needles from SO<sub>2</sub> fumigated and control seedlings were excised into 1 cm sections and transferred to a chilled solution (1:10, W/V) containing either 0.05 M Tris-HCl buffer or 0.1 N sodium phosphate buffer (pH 7.4) and 1% PVP (polyvinyl pyrrolidone) and 1 mM DTE (dithioerythritol). The tissue was homogenized at moderate speed for 1.5 min by a Brinkman Polytron homogenizer. The homogenate was passed through two layers of cheese cloth and the filtrate was immediately used as "crude enzyme extract".

## 2.8 ENZYME ASSAYS

### 2.8.1 Acid Phosphatase

A typical reaction mixture contained 0.6 mM p-nitrophenyl phosphate (P-NPP), 0.05 M acetate buffer (pH 5.0) and 25  $\mu$ l of "crude enzyme extract" (from SO<sub>2</sub> fumigated plant material) in a final volume of 2 ml. The reaction was run for 30 min in a water bath at 30°C and stopped by the addition of 2 ml of 10% Na<sub>2</sub>CO<sub>3</sub>. A control in which P-NPP was added after the addition of Na<sub>2</sub>CO<sub>3</sub> was always run with the regular reaction set. This mixture was clarified by centrifugation and the acid phosphatase activity was determined by measuring the difference in optical density between the control and the reaction solution at 410 nm.

### 2.8.2 Malate Dehydrogenase

The enzyme activity was assayed by following the reverse reaction (L-malate + NAD  $\rightleftharpoons$  Oxaloacetate + NADH + H<sup>+</sup>) as

described by Ochoa (1955). A typical assay mixture included 0.3 ml of 0.0015M NADH, 0.1 ml of 0.1 M KCN, 0.1 ml of 0.0076M oxaloacetate (cis), 2.5 ml of 0.1 M sodium phosphate buffer (pH 7.5) and 10  $\mu$ l of enzyme extract. The rate of the reaction was monitored by measuring the decrease in absorbance at 340 nm against a blank containing the reaction mixture without NADH.

## 2.9 LIPID BIOSYNTHESIS

Excised tissue (0.5 g) from fumigated and control jack pine seedlings was transferred to flasks containing 10 ml water and 10  $\mu$ Ci acetate [ $1^{-14}$ C]. The incubations were carried out for 3 h in a gyrotary water bath shaker at 30°C and under 200  $\mu$ E  $m^{-2}s^{-1}$  light intensity. After incubation, each tissue sample was removed, washed with distilled water and homogenized in 50 ml (chloroform-methanol (2:1)). The homogenate was then filtered through a glass-wool-packed Buchner funnel, and the residue was washed repeatedly with chloroform-methanol. The chloroform extracts containing lipids were washed with acidic water and evaporated to dryness under reduced pressure and the labelled products analyzed by thin-layer-chromatography.

Mention of a trademark does not imply its approval to the exclusion of other products or vendors that may also be suitable.

### 3. RESULTS AND DISCUSSION

The experiments for each study were conducted several times and the trends for SO<sub>2</sub> response were found to be highly reproducible. Although considerable effort was made to select uniformly grown plant material, it was difficult to eliminate the biological variability on a day-to-day basis. In order to overcome this difficulty each experiment was designed to include a control (non SO<sub>2</sub>-treated plant material) which provided meaningful data on SO<sub>2</sub> response.

#### 3.1 EFFECTS OF AQUEOUS SO<sub>2</sub> ON TOTAL CHLOROPHYLL CONTENT

The pine needles that were incubated in 10, 25, 50 and 100 ppm aqueous SO<sub>2</sub> remained green and looked quite normal whereas the higher concentrations caused increasing degrees of discoloration or bleaching.

The low concentrations (10-100 ppm SO<sub>2</sub>) did not produce any appreciable change in total chlorophyll content. However, when SO<sub>2</sub> concentration was increased from 100 ppm to 250 and 550 ppm, there was a sharp drop in pH of the incubation medium as well as in total chlorophyll content (Table 1). At 500 ppm, the pH dropped to 3.95 and the chlorophyll content was reduced to about half that of the control values. From these results it would appear that the decrease in chlorophyll content could be attributed to a drop in pH. The HCl control adjusted to the same pH as produced by 500 ppm SO<sub>2</sub>, however, showed approximately 4% decrease in chlorophyll content compared with the regular control. It is therefore evident that the pigment destruction resulted from direct effects of SO<sub>2</sub> in addition to indirect effects such as increase in acidity.

##### 3.1.1 Chlorophyll Breakdown

In order to understand the mechanism of chlorophyll destruction, the effect of aqueous SO<sub>2</sub> on Mg<sup>++</sup> loss from

Table 1. The effect of various concentrations of  $\text{SO}_2$  on total chlorophyll content of pine needles.

Treatment	pH of Incubation Medium	Chlorophyll Content (mg/l)	% of Control
Control	7.20	65.27	100.00
10 ppm $\text{SO}_2$	7.18	64.13	98.25
25 ppm $\text{SO}_2$	7.13	63.50	97.29
50 ppm $\text{SO}_2$	7.10	63.82	97.78
100 ppm $\text{SO}_2$	7.00	63.38	97.10
250 ppm $\text{SO}_2$	6.60	47.98	73.51
500 ppm $\text{SO}_2$	3.95	34.69	53.15
HCl Control	3.95	62.71	96.08



chlorophyll (conversion of chlorophyll into phaeophytin) was determined. Some of the mechanisms of chlorophyll breakdown are shown in Figure 3. Acids such as HF and HCl have been shown to convert chlorophyll into phaeophytin with the release of  $Mg^{++}$  (Rao and Le Blanc 1965; Oker 1967).  $Mg^{++}$  is replaced by 2 molecules of hydrogen with a resulting change in the light absorption spectral properties of chlorophyll molecules.

Table 2 shows the release of  $Mg^{++}$  from excised pine needles as a result of treatment with high concentration of aqueous  $SO_2$ . The loss of  $Mg^{++}$  from needles treated with lower concentrations of  $SO_2$  (10-50 ppm) was minimal (about 7% at 50 ppm  $SO_2$ ). However, such loss increased considerably with increasing  $SO_2$  concentration. The data also show that even at very low pH, the conversion of chlorophyll into phaeophytin is due primarily to  $SO_2$  and only to a small extent to acidity.

### 3.1.2 Effect on Various Pigments

The data in Figure 4 show changes in the content of chlorophyll a (A), chlorophyll b (B), chlorophyllide a (C), chlorophyllide b (D), and phaeophytin a (E) and phaeophytin b (F) as affected by various concentrations of aqueous  $SO_2$ . The amounts of chlorophyll a and chlorophyll b declined with increasing concentrations of aqueous  $SO_2$ . However, chlorophyll a is more sensitive to  $SO_2$  attack than chlorophyll b. At 500 ppm  $SO_2$  chlorophyll a was almost completely destroyed, but only one-half of chlorophyll b was destroyed.

Low concentrations of  $SO_2$  (10-100 ppm) did not have any appreciable effect on phaeophytin a (E) content but the content of this pigment increased considerably at 250 and 500 ppm. Phaeophytin b was absent in the control tissue and its content remained unchanged with increasing  $SO_2$ , suggesting conversion of only chlorophyll a to phaeophytin a. Since no chlorophyllide a (C) was detected in the control tissue and  $SO_2$  did not show any effect on its content, it is suggested that




Figure 3. Different mechanisms of chlorophyll breakdown.

Table 2. The effect of various concentrations of aqueous  $\text{SO}_2$  on  $\text{Mg}^{++}$  loss from pine needle chlorophyll.

Treatment	pH of Incubation Medium	$\text{Mg}^{++}$ Loss (mg/l)	% of Control
Control	7.20	7.94	100.00
10 ppm $\text{SO}_2$	7.18	8.07	101.64
25 ppm $\text{SO}_2$	7.13	8.26	104.03
50 ppm $\text{SO}_2$	7.10	8.37	105.42
100 ppm $\text{SO}_2$	7.00	9.05	113.98
250 ppm $\text{SO}_2$	6.60	26.81	337.66
500 ppm $\text{SO}_2$	3.95	32.76	412.59
HCl Control	3.95	9.73	112.54




Figure 4. The effect of aqueous SO<sub>2</sub> on chlorophyll a (A), chlorophyll b (B), chlorophyllide a (C), chlorophyllide b (D), phaeophytin a (E) and phaeophytin b (F).

chlorophyll a was destroyed by processes other than simple transformations. The amount of chlorophyllide b (D) increased with increasing concentration of  $\text{SO}_2$  to a maximum of 50 ppm followed by a drop at 100 ppm that was sustained up to 500 ppm  $\text{SO}_2$ . The increase in the amount of chlorophyllide b indicates that at least a part of the chlorophyll b was converted into chlorophyllide b.

### 3.1.3 Effect on Chlorophyllase Activity

Figure 4 indicates that part of the chlorophyll b was converted into chlorophyllide b and this conversion was maximum at 50 ppm  $\text{SO}_2$ . Since chlorophyllase is capable of such conversion it was thought desirable to determine if  $\text{SO}_2$  has any effect on the activity of pine chlorophyllase. Chlorophyllase activity was measured in terms of chlorophyllide formation.

Concentrations of 10-50 ppm  $\text{SO}_2$  stimulated the chlorophyllase activity while higher concentrations (100-500 ppm) inhibited it (Figure 5). Maximum stimulation was obtained at 50 ppm. The activity dropped to a value close to that of control at 100 ppm and then exhibited significant inhibition with increasing concentrations of  $\text{SO}_2$ . Maximum inhibition (about 30%) was obtained with 500 ppm  $\text{SO}_2$ . This inhibition was at first attributed to the low pH of the incubation medium caused by high concentration of  $\text{SO}_2$ , but HCl control showed inhibition of less than 10%. Therefore, the inhibition of chlorophyllase activity at high concentrations of  $\text{SO}_2$  is not entirely due to pH effect. It appears that  $\text{SO}_2$  has a direct effect on chlorophyllase activity.

These results clearly indicate that the chlorophyllase activity index is a very sensitive indicator of  $\text{SO}_2$  concentrations. This method may be useful for detecting  $\text{SO}_2$  pollution before the development of visual symptoms.

These biochemical studies suggest that low concentrations of  $\text{SO}_2$  which normally do not produce any visual symptoms on the foliage can have an effect on specific enzyme systems such as chlorophyllase. Since low concentrations of  $\text{SO}_2$  were shown to




Figure 5. The effect of aqueous  $\text{SO}_2$  on the activity of chlorophyllase.

have no appreciable effects on  $Mg^{++}$  loss, injury may be a result of increased activity of chlorophyllase. On the other hand, high concentrations of  $SO_2$  that are capable of producing visual symptoms may cause injury by inhibiting the activities of important enzyme systems, by destroying chlorophyll pigments, and thereby resulting in a net drop in photosynthetic efficiency.

### 3.2 ULTRASTRUCTURAL STUDIES

Electron microscopy has been used to demonstrate the extent of ultrastructural injury caused by air pollutants such as peroxyacetyl nitrate and ozone (Thompson *et al.* 1965, 1966). Little is known about  $SO_2$  effects on ultrastructural organization and metabolic activities in pine tissues. Wellburn *et al.* (1972), however, reported the effects of  $SO_2$  on the ultrastructure of broad bean chloroplasts.

According to cytological examinations, the tip of pine needles is composed of old and fully differentiated cells and the base contains very young and actively growing cells (Linzon, 1967). About 1 cm sections excised from the tip and base of pine needles were incubated in different concentrations of aqueous  $SO_2$  solutions and the ultrastructural changes in chloroplasts were examined by means of electron microscopy.

At aqueous concentrations of 100 and 500 ppm,  $SO_2$  caused swelling of thylakoid discs and desintegrated other intrachloroplast membranes, resulting in the formation of small vesicles (in older, matured tissue). Chloroplast structural injury was more pronounced in old tissues than in younger and metabolically active tissues. In general, electron microscopy showed that older, fully matured tissue was more sensitive to  $SO_2$  injury than the younger actively growing tissue. This was shown by disorientation of chloroplast structure, dislocation of chloroplasts and absence of mitochondria in the older tissue treated with high concentrations of  $SO_2$ . The detrimental effects of  $SO_2$  on the ultrastructural organization of pine needle cells appeared to be

markedly different from the effects of low pH alone (HCl control).

Possible relationship between structural disorganization and disruption of biochemical activities of chloroplasts from SO<sub>2</sub>-treated young and old tissues were determined by measuring the photosynthetic activity such as Hill reaction (Malhotra 1976) and <sup>14</sup>CO<sub>2</sub> fixation (Malhotra 1977) of isolated chloroplasts. Biochemical observations (photosynthetic activity) were in good agreement with the cytological observations.

### 3.3 EFFECT OF SO<sub>2</sub> ON TISSUE GLYCOLIPIDS AND PERMEABILITY PROPERTIES

Chloroplast membranes are sites of many enzymatic reactions and are made up of two very important structural components namely lipids and proteins. It is possible that structural disruption due to SO<sub>2</sub> has been preceded by changes in a number of biochemical events.

Chloroplast lipids are mostly glycolipids namely monogalactosyldiglycerides (MGDG), digalactosyldiglycerides (DGDG) and sulphoquinovosyldiglycerides (SQDG) (James and Nichols, 1966). These lipids are known to be involved in the structure and function of chloroplast membranes (Shaw *et al.* 1976; Erwin and Bloch 1963); however, nothing is known about the effect of SO<sub>2</sub> on metabolism of these compounds. We therefore decided to study the effect of aqueous SO<sub>2</sub> on (1) the level of these lipids in pine needles and (2) permeability properties of pine needle tissue.

The data in Figure 6 show changes in the levels of glycolipids from fully developed and young needles as affected by various concentrations of aqueous SO<sub>2</sub>. Concentrations of MGDG, DGDG and SQDG declined sharply with increasing concentrations of SO<sub>2</sub> up to 25 ppm. Further increase in SO<sub>2</sub> concentration (100 ppm) resulted in only small changes in the concentration of these glycolipids. An interesting observation is that the effect of SO<sub>2</sub> on the level of SQDG was more pronounced in the fully






Figure 6. The effect of aqueous  $\text{SO}_2$  on the levels of glycolipids in fully developed and young pine needles.

developed tissues than in the young tissues. The reverse was true of the DGDG fraction. These differences clearly indicate that the biosensitivity of various cellular metabolites to  $\text{SO}_2$  depends on the age of the needles. It is also interesting to note that a dramatic decrease in the level of all these lipids was obtained at a very low concentration of  $\text{SO}_2$  (50 ppm). Such low concentrations did not produce any visual symptoms on pine needles.

The influence of aqueous  $\text{SO}_2$  on cellular metabolites such as soluble sugars was determined to ascertain whether the effects of  $\text{SO}_2$  discussed were confined to the membrane glycolipids or whether other cellular metabolites such as sugars were also influenced. The effect of  $\text{SO}_2$  on the release of sugars would indicate changes in cellular permeability. The incubation of tissues from both fully developed and younger needles resulted in a release of soluble sugars from the tissues, even in the absence of  $\text{SO}_2$  (Figure 7). However, the presence of  $\text{SO}_2$  in the incubation solution caused a dramatic increase in the amount of sugars released from both the tissues. The release of soluble sugars from tissues of both ages might have occurred due to the decrease in glycolipid concentrations in the  $\text{SO}_2$ -treated tissues. However, the amount of sugars released from the young tissues was considerably higher than the glycolipid losses, suggesting that  $\text{SO}_2$  caused a release of some other bound form(s) of sugar.

Sulphur dioxide was also shown to have an inhibitory effect on other essential plant metabolites such as amino acids and organic acids. The lack of these metabolites can impair important processes such as photosynthesis and respiration.

The difference in biochemical responses to aqueous  $\text{SO}_2$  between the fully developed and young pine needle tissues indicates that the phytotoxicity of  $\text{SO}_2$  is at least partially dependent upon the stage of plant tissue development. In view of the importance of galactolipids for the structural integrity of the chloroplast thylakoids, any changes in these lipids caused by  $\text{SO}_2$  would result in structural and functional injuries to the




Figure 7. The effect of aqueous  $\text{SO}_2$  on the release of soluble sugars from fully developed and young pine needles.

biomembranes. We feel that the process of hidden  $\text{SO}_2$  injuries such as swelling of thylakoid discs and disintegration of intrachloroplast membranes (Malhotra 1976) may be initiated as a result of alteration in glycolipid components. Since very low concentrations of aqueous  $\text{SO}_2$  (10-25 ppm) usually do not produce any visual symptoms but cause marked changes in the glycolipid content and composition, the analysis of such lipids may provide an effective tool for determining the extent of hidden  $\text{SO}_2$  injury to vegetation.

#### 3.4 EFFECTS OF GASEOUS $\text{SO}_2$ ON PHOSPHATASE ACTIVITY

Under field conditions, vegetation is normally exposed to air pollutants on an intermittent basis. This allows the vegetation to recover at least partially from the air pollutant stress. In order to simulate such field conditions, the plants were fumigated with various  $\text{SO}_2$  concentrations for a known length of time and then placed in a  $\text{SO}_2$ -free environment to determine the percent recovery of impaired biochemical functions.

The results in Table 3 show that exposure of pine seedlings at 0.35 ppm  $\text{SO}_2$  for 24 h produced a marked inhibition in acid phosphatase activity in both young (top clustre of new growth) and old (fully developed, approximately 3 month old) needles. Enzyme activity was inhibited more in the older needles than in the younger ones. Inhibition of enzyme activity further increased during the 24 h recovery period in a  $\text{SO}_2$ -free environment. There was very little visual symptom development after the 24 h fumigation. However, the symptoms were much more obvious during the next 24 h recovery period indicating a lag period requirement for such symptom development. These results, therefore, suggest that fumigation of pine seedlings under above conditions may result in serious impariment of plant metabolism. Rabe and Kreeb (1976) and Schmid and Kreeb (1975) have reported a similar inhibition of acid phosphatase activity in alfalfa plants and lichens after treatment of  $\text{SO}_2$ .

Table 3. The effects of gaseous SO<sub>2</sub> on acid phosphatase activity in young and old needles of jack pine seedlings and its ability to recover in a SO<sub>2</sub>-free atmosphere.

Conc. (ppm)	Fumigation Conditions			% of Control Activity	
	Duration (h)	SO <sub>2</sub> Uptake/ tree/h (ppb)	Recovery Period (h)	Young Needles <sup>a</sup>	Old Needles <sup>a</sup>
Expt. 1					
0.35	24	20	0	64	46
0.35	24	20	24	54	36
Expt. 2					
0.33	1	9	0	90	82
0.33	1	9	24	100	100
Expt. 3					
0.10	24	4	0	80	75

<sup>a</sup> Young and old needles represented the pale green top clustre and the dark green 3-4 month old growth, respectively.

The fumigation of pine seedlings at 0.33 ppm SO<sub>2</sub> for only 1 h also inhibited enzyme activity in both young and old tissues by about 10% (Table 3, Expt. 2). However, this inhibition was completely recovered after 24 h in a SO<sub>2</sub>-free atmosphere. In this experiment, the treatment of plants with SO<sub>2</sub> resulted in an inhibition of phosphatase activity without development of any visual symptoms of SO<sub>2</sub> phytotoxicity.

Plants fumigated with 0.1 ppm SO<sub>2</sub> for 24 h inhibited the enzyme activity in young needles by 20% and in old needles by 25% (Table 3, Expt. 3). These plants did not produce visual symptoms of SO<sub>2</sub> phytotoxicity. The results clearly suggest that SO<sub>2</sub> when applied at either low concentration for a long period or high concentration for a short period, can inhibit the acid phosphatase activity in jack pine seedlings. This inhibition can become more severe with increasing SO<sub>2</sub> concentration or increasing exposure time.

Table 3 also shows that the effect of SO<sub>2</sub> on acid phosphatase was dependent upon the amount of SO<sub>2</sub> taken up per seedling. The variability in the amount of SO<sub>2</sub> uptake per seedling per hour between Expt. 1 and Expt. 2 (Table 3) was a result of low humidity conditions in the later case. It is suggested that stomata in Expt. 2 were probably only partially open resulting in considerably less SO<sub>2</sub> uptake per seedling.

Our results suggest that the acid phosphatase activity could be used as a very sensitive biochemical indicator of SO<sub>2</sub> phytotoxicity. The ability of the SO<sub>2</sub>-fumigated plants to recover their enzymatic activity in a SO<sub>2</sub>-free environment suggests a transient nature of SO<sub>2</sub> effect. However, irreversible injury can result at high SO<sub>2</sub> concentrations (Table 3, Expt. 1).

### 3.5 EFFECT OF GASEOUS SO<sub>2</sub> ON MALATE DEHYDROGENASE

The results in Table 4 demonstrate an inhibitory effect of SO<sub>2</sub> on malate dehydrogenase. The activity of this enzyme declined in relation to the length of fumigation, reaching

Table 4. The effects of SO<sub>2</sub> (0.34 ppm) on malate dehydrogenase activity from pine needles.

Length of Fumigation (h)	Control	Malate Dehydrogenase Activity	% Inhibition
		(units <sup>a</sup> /mg protein)	
19	303	274	10
72	362	226	37
92	315	164	48

<sup>a</sup>One unit is defined as the amount of enzyme required to cause an O.D change of 0.01/min at 340 mμ.

a value almost 50 percent of the control value after 92 h. It is suggested that  $\text{SO}_2$  affects the plant metabolism probably by interfering with important enzyme systems such as malate dehydrogenase. The related metabolites such as organic and amino acids also showed a substantial decline after fumigation of jack pine seedlings with 0.34 and 0.51 ppm  $\text{SO}_2$ .

Malate dehydrogenase assay also has potential as a reliable assessment tool for  $\text{SO}_2$  phytotoxicity prior to visual symptom development.

### 3.6 EFFECT OF GASEOUS $\text{SO}_2$ ON LIPID BIOSYNTHESIS

Exposure of jack pine seedlings for 24 h to 0.18-0.20 ppm  $\text{SO}_2$  markedly inhibited the biosynthesis of various lipids (Table 5, Expt. 1, 2). The inhibitory effect among the phospholipids was not specifically greater on any one component. Among the galactolipids, DGDG was generally inhibited more than MGDG. Fumigation of seedlings for 24 h with a high  $\text{SO}_2$  concentration (0.35 ppm) drastically inhibited all lipid biosynthesis (Table 5, Expt. 1) and caused needle tip burn. It is therefore suggested that invisible and visible effects of prolonged  $\text{SO}_2$  exposure are directly related to impaired cellular metabolism.

#### 3.6.1 Recovery of Fumigated Seedlings

In order to simulate the field conditions, such as varying concentrations of  $\text{SO}_2$ , as closely as possible, fumigations were carried out in controlled environment cuvettes and followed by different lengths of recovery time in a  $\text{SO}_2$ -free environment. The experimental conditions chosen and the results obtained are given in Table 5 (Expt. 2, 3). A high  $\text{SO}_2$  concentration (0.37 ppm) for a short exposure time (1 h) resulted in much more inhibition of lipid biosynthesis than a low  $\text{SO}_2$  concentration (0.20 ppm) for a long period of time (24 h) (Table 5, Expt. 2, 3). The inhibitory effects of either fumigation were partially or completely reversed upon removal of plants from the



Table 5. The effects of gaseous SO<sub>2</sub> on lipid biosynthesis in developing needles from jack pine seedlings and their ability to recover in a SO<sub>2</sub>-free environment.

Fumigation Conditions		Recovery Period (h)	% of Control							
Conc. (ppm)	Duration (h)		PC <sup>a</sup>	PE <sup>a</sup>	PG <sup>a</sup>	DGDG <sup>a</sup>	MGDG <sup>a</sup>	DG <sup>a</sup>	FFA <sup>a</sup>	TG <sup>a</sup>
Expt 1										
0.18	24	0	78.2	74.0	68.6	57.4	81.1	75.6	75.0	80.4
0.35	24	0	30.0	44.0	41.6	29.8	26.4	26.7	15.6	39.0
Expt 2										
0.20	24	0	81.8	60.5	61.5	73.1	81.1	80.6	76.5	88.9
0.20	24	48	82.2	70.3	67.4	100.0	100.0	100.0	100.0	100.0
Expt 3										
0.37	1	0	43.6	37.7	33.3	42.3	44.8	40.7	42.2	39.3
0.37	1	24	63.4	65.4	69.4	66.0	59.2	63.6	115.0	52.4

<sup>a</sup>PC, phosphatidyl choline; PE, phosphatidyl ethanolamine; PG, phosphatidyl glycerol; DGDG, digalactosyl-diglycerides; MGDG, monogalactosyldiglycerides; DG, diglycerides; FFA, free fatty acids; and TG, triglycerides.

SO<sub>2</sub> environment. Recovery of biosynthetic ability of plants exposed to a high SO<sub>2</sub> concentration (0.37 ppm) for the short duration (1 h), however, was slower than that of the plants exposed to low SO<sub>2</sub> (0.20 ppm) for a longer period (24 h). It is also interesting to note that at high SO<sub>2</sub> concentration (0.35-0.37 ppm) much of the damage to lipid biosynthesis of tissues occurred during the initial period of exposure (Table 5, Expt. 1 vs Expt. 3). The ability of plants to recover their synthetic activities after removal of SO<sub>2</sub> from the environment is of great significance; however, the recovery is dependent upon the extent of SO<sub>2</sub> exposure, magnitude of metabolic injury, and length of recovery time.

### 3.7 BIOCHEMICAL ANALYSES ON FIELD SAMPLES

Assessment of air pollution injury to jack pine and white spruce in the impingement area south of GCOS plant (Figure 2) was carried out using some of the more sensitive biochemical and physiological techniques developed in our laboratory. Branches from the two species were collected in a gradient at a distance of 2.8, 4.2 and 8.3 km south of the GCOS plant and analyzed for sulphur content, CO<sub>2</sub> gas exchange capacity (under controlled temperature and light conditions), chlorophyll/phaeophytin ratios, and activity of highly sensitive enzymes such as phosphatase and malate dehydrogenase. Although a gradient in sulphur content was evident, there was no difference in any of these biochemical and physiological functions that responded to SO<sub>2</sub> under laboratory conditions. The highest sulphur levels observed in jack pine needles from the above sites were close to the lower end of the normal sulphur range for such tissues. Low sulphur content and the transient nature of SO<sub>2</sub> injury (demonstrated under laboratory conditions) may be the reasons for not being able to detect SO<sub>2</sub> response in the field collected samples.

The work conducted so far, under controlled conditions, suggested that low concentrations of SO<sub>2</sub> which normally do not

produce any visual symptoms on the foliage can cause injury to forest vegetation by inhibiting lipid synthesis and activities of various important enzyme systems.

#### 4. CONCLUSIONS

Biochemical and ultrastructural responses to aqueous and gaseous  $\text{SO}_2$  show that, depending upon the concentration and length of exposure time,  $\text{SO}_2$  can cause considerable injury to forest species prior to visual symptom development. However, such injury can either be partially or fully recovered in a  $\text{SO}_2$ -free environment. The techniques developed to date have great potential as diagnostic tools to: (1) detect  $\text{SO}_2$  injury to vegetation before the development of visual symptoms of  $\text{SO}_2$  phytotoxicity, (2) establish threshold levels of air pollutant injury to vegetation and (3) screen candidate revegetation species on the basis of their true tolerance to air pollutants.

## 5. REFERENCES CITED

- Barrie, L.A., and D.M. Whelpdale. 1978. Background air and precipitation chemistry. Pages 124-159 in F. Fanaki, comp. Meteorology and air quality winter field study in the AOSERP study area, March 1976. Prep. for the Alberta Oil Sands Environmental Research Program by Fisheries and Environment Canada, Atmospheric Environment Service. AOSERP Report 27. 249 pp.
- Bleasdale, J.K.A. 1952. Atmospheric pollution and plant growth. *Nature* 169:376-377.
- Brandt, C.S., and W.W. Heck. 1968. Effects of air pollutants on vegetation. Volume 1, Pages 401-443 in A.C. Stern, ed. *Air Pollution*. Academic Press, N.Y. 694 pp.
- Coker, P.D. 1967. The effects of SO<sub>2</sub> pollution on bark epiphytes. *Trans. Br. Bryol. Soc.* 5:341-347.
- Dubois, M., K.A. Gilles, J.K. Hamilton, P.A. Rebers, and F. Smith. 1956. Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 28:350-356.
- Erwin, J., and K. Bloch. 1963. Polyunsaturated fatty acid in some photosynthetic micro-organisms. *Biochem. Z.* 338-496.
- Folch, J., M. Lees, and G.H. Sloane-Stanley. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J. Biol. Chem.* 226:497-509.
- Hocking, D., and M.B. Hocking. 1977. Equilibrium solubility of trace atmospheric SO<sub>2</sub> in water and its bearing on air pollution injury to plants. *Environ. Pollut.* 13:57-64.
- Holden, M. 1961. The breakdown of chlorophyll by chlorophyllase. *Biochem. J.* 78:359-364.
- James, A.T., and B.W. Nichols. 1966. Lipids of photosynthetic systems. *Nature* 210:372-375.
- Linzon, S.N. 1967. Histological studies of symptoms in semi-mature tissue needle blight of eastern white pine. *Can. J. Bot.* 45:133-143.
- Malhotra, S.S. 1976. Effects of SO<sub>2</sub> on biochemical activity and ultrastructural organization of pine needle chloroplasts. *New Phytol.* 76:239-245.

- Malhotra, S.S. 1977. Effects of aqueous SO<sub>2</sub> on chlorophyll destruction in *Pinus contorta*. New Phytol. 78:101-109.
- Malhotra, S.S., and D. Hocking. 1976. Biochemical and cytological effects of sulphur dioxide on plant metabolism. New Phytol. 76:227-237.
- Ochoa, S. 1955. Malate dehydrogenase from pig heart. Volume 1, Pages 735-739 in S.P. Colowick and N.O. Kaplan, eds. Methods in enzymology. Academic Press, N.Y. 835 pp.
- Rabe, S., and K. Kreeb. 1976. Ein Methode zur Laborbegasung von test Planzen mit Schwefeldioxide and ihre Anwendung bei Untersuchungen zur Enzymaktivitat. Angew Bot. 50:71-78.
- Rao, D.N., and F. LeBlanc. 1965. Effects of SO<sub>2</sub> on the lichen algae with special reference to chlorophyll. The Bryologist 69:69-75.
- Roughan, P.G., and R.D. Batt. 1968. Quantitative analysis of sulpholipid (sulphoquinovosyl diglyceride) and galactolipids (monogalactosyl digalactosyl diglycerides) in plant tissue. Anal. Biochem. 22:74-88.
- Saunders, P.J.W. 1966. The toxicity of SO<sub>2</sub> to *Diplocarpon rosae* Wolf causing black spot of roses. Ann. Appl. Biol. 58:103-114.
- Saunders, P.J.W., and C.M. Wood. 1973. SO<sub>2</sub> in the environment, its production dispersal and fate. Pages 7-37 in B.W. Ferry, M.S. Baddeley and D.L. Hawksworth, eds. Air Pollution and lichens. University of Toronto Press, Toronto. 389 pp.
- Schmid, M.L., and K. Kreeb. 1975. Enzymatische indikation gas-geschaedigter Flechten. Angew. Bot. 49:141-154.
- Shaw, A.B., M.M. Anderson, and R.E. MacCarty. 1976. Role of galactolipids in spinach chloroplast lamellar membranes. II. Effect of galactolipid depletion in phosphorylation and electron flow. Plant Physiol. 57:724-729.
- Thompson, W.W., W.M. Dugger, and R.L. Palmer. 1965. Effects of peroxyacetyl nitrate on ultrastructure of chloroplasts. Bot. Gaz. 126:66.
- Thompson, W.W., W.M. Dugger, and R.L. Palmer. 1966. Effects of ozone on the fine structure of the palisade parenchyma cells of bean leaves. Can. J. Bot. 44:1677-1682.

- Tingey, D.T., W.W. Heck, and R.A. Reinert. 1971. Effect of low concentration of  $O_3$  and  $SO_2$  on foliage, growth and yield of radish. J. Am. Soc. Hort. Sci. 96:369-371.
- Wellburn, A.R., Majernik, O., and F.A.M. Wellburn. 1972. Effects of  $SO_2$  and  $NO_2$  polluted air upon the ultrastructure of chloroplasts. Environ. Pollut. 3:37-49.
- White, R.C., I.D. Jones, and E. Gibbs. 1963. Determination of chlorophylls, chlorophyllides, phaeophytins and phaeophorbides in plant material. J. Food Sci. 28:431-436.
- Ziegler, I. 1975. The effect of  $SO_2$  pollution on plant metabolism. Residue Rev. 56:79-105.

## 6. AOSERP RESEARCH REPORTS

1. AOSERP First Annual Report, 1975
2. AF 4.1.1 Walleye and Goldeye Fisheries Investigations in the Peace-Athabasca Delta--1975
3. HE 1.1.1 Structure of a Traditional Baseline Data System
4. VE 2.2 A Preliminary Vegetation Survey of the Alberta Oil Sands Environmental Research Program Study Area
5. HY 3.1 The Evaluation of Wastewaters from an Oil Sand Extraction Plant
6. Housing for the North--The Stackwall System
7. AF 3.1.1 A Synopsis of the Physical and Biological Limnology and Fisheries Programs within the Alberta Oil Sands Area
8. AF 1.2.1 The Impact of Saline Waters upon Freshwater Biota (A Literature Review and Bibliography)
9. ME 3.3 Preliminary Investigations into the Magnitude of Fog Occurrence and Associated Problems in the Oil Sands Area
10. HE 2.1 Development of a Research Design Related to Archaeological Studies in the Athabasca Oil Sands Area
11. AF 2.2.1 Life Cycles of Some Common Aquatic Insects of the Athabasca River, Alberta
12. ME 1.7 Very High Resolution Meteorological Satellite Study of Oil Sands Weather: "a Feasibility Study"
13. ME 2.3.1 Plume Dispersion Measurements from an Oil Sands Extraction Plant, March 1976
14. HE 2.4 Athabasca Oil Sands Historical Research Project. Volume I: Design
15. ME 3.4 A Climatology of Low Level Air Trajectories in the Alberta Oil Sands Area
16. ME 1.6 The Feasibility of a Weather Radar near Fort McMurray, Alberta
17. AF 2.1.1 A Survey of Baseline Levels of Contaminants in Aquatic Biota of the AOSERP Study Area
18. HY 1.1 Interim Compilation of Stream Gauging Data to December 1976 for the Alberta Oil Sands Environmental Research Program
19. ME 4.1 Calculations of Annual Averaged Sulphur Dioxide Concentrations at Ground Level in the AOSERP Study Area
20. HY 3.1.1 Characterization of Organic Constituents in Waters and Wastewaters of the Athabasca Oil Sands Mining Area



21. AOSERP Second Annual Report, 1976-77
22. HE 2.3 Maximization of Technical Training and Involvement of Area Manpower
23. AF 1.1.2 Acute Lethality of Mine Depressurization Water on Trout Perch and Rainbow Trout
24. ME 4.2.1 Review of Dispersion Models and Possible Applications in the Alberta Oil Sands Area
25. ME 3.5.1 Review of Pollutant Transformation Processes Relevant to the Alberta Oil Sands Area
26. AF 4.5.1 Interim Report on an Intensive Study of the Fish Fauna of the Muskeg River Watershed of Northeastern Alberta
27. ME 1.5.1 Meteorology and Air Quality Winter Field Study in the AOSERP Study Area, March 1976
28. VE 2.1 Interim Report on a Soils Inventory in the Athabasca Oil Sands Area
29. ME 2.2 An Inventory System for Atmospheric Emissions in the AOSERP Study Area
30. ME 2.1 Ambient Air Quality in the AOSERP Study Area, 1977
31. VE 2.3 Ecological Habitat Mapping of the AOSERP Study Area: Phase I
32. AOSERP Third Annual Report, 1977-78
33. TF 1.2 Relationships Between Habitats, Forages, and Carrying Capacity of Moose Range in northern Alberta. Part I: Moose Preferences for Habitat Strata and Forages.
34. HY 2.4 Heavy Metals in Bottom Sediments of the Mainstem Athabasca River System in the AOSERP Study Area
35. AF 4.9.1 The Effects of Sedimentation on the Aquatic Biota
36. AF 4.8.1 Fall Fisheries Investigations in the Athabasca and Clearwater Rivers Upstream of Fort McMurray: Volume I
37. HE 2.2.2 Community Studies: Fort McMurray, Anzac, Fort MacKay
38. VE 7.1.1 Techniques for the Control of Small Mammals: A Review
39. ME 1.0 The Climatology of the Alberta Oil Sands Environmental Research Program Study Area
40. VE 7.1 Interim Report on Reclamation for Afforestation by Suitable Native and Introduced Tree and Shrub Species
41. AF 3.5.1 Acute and Chronic Toxicity of Vanadium to Fish
42. TF 1.1.4 Analysis of Fish Production Records for Registered Traplines in the AOSERP Study Area, 1970-75
43. TF 6.1 A Socioeconomic Evaluation of the Recreational Fish and Wildlife Resources in Alberta, with Particular Reference to the AOSERP Study Area. Volume I: Summary and Conclusions
44. VE 3.1 Interim Report on Symptomology and Threshold Levels of Air Pollutant Injury to Vegetation, 1975 to 1978
45. VE 3.3 Interim Report on Physiology and Mechanisms of Air-Borne Pollutant Injury to Vegetation, 1975 to 1978

46. VE 3.4 Interim Report on Ecological Benchmarking and  
Biomonitoring for Detection of Air-Borne Pollutant  
Effects on Vegetation and Soils, 1975 to 1978

These reports are not available upon request. For further information about  
availability and location of depositories, please contact:

Alberta Oil Sands Environmental Research Program  
15th Floor, Oxbridge Place  
9820-106 Street  
Edmonton, Alberta T5K 2J6