

**VALIDATION OF BASAL DIAMETER RATIO
COMPETITION INDEX FOR LODGEPOLE
PINE-ASPEN**

**SECOND YEAR POST-TREATMENT
PROGRESS REPORT**

D.A. MacIsaac
Canadian Forest Service
Edmonton, Alberta

1996

DISCLAIMER

The project/study on which this report is based was funded by the Foothills Model Forest under the Partners in Sustainable Development of Forests initiative delivered by the Canadian Forest Service of Natural Resources Canada and funded by Canada's Green Plan for a Healthy Environment.

The views, statements and conclusions expressed and the recommendations made in this report are entirely those of the authors and should not be construed as the statements or conclusions of, or as expressing the opinions of, Foothills Model Forest.

FOOTHILLS MODEL FOREST MISSION

"to develop and recommend an approach to sustainability and integrated resource management through research and technology developed by means of collaborative partnerships".

RELATIONSHIP BETWEEN FOOTHILLS MODEL FOREST AND RESOURCE MANAGEMENT AGENCIES

The Foothills Model Forest represents a broad range of stakeholder groups with interest in Alberta's forests and how they are managed. However, Foothills Model Forest has no resource management authority or responsibility. The authority over, and responsibility for, the management of Alberta's public lands is vested in the Government of Alberta. The Government delegates certain rights and responsibilities to various resource industries and organizations which conduct their activities on public lands in Alberta. The Government of Alberta and other agencies and organizations will consider and respond to the recommendations of Foothills Model Forest from the perspective of their particular rights, responsibilities, obligations and stewardship commitments.

ABSTRACT

In 1993, a four year Foothills Model Forest study was initiated to verify the use of the recently-developed Basal Diameter Ratio Competition Index in stand tending decisions for juvenile lodgepole pine-aspen competition in west-central Alberta. A mixed-nested experiment with three blocks and four levels of aspen removal (treatments) was designed. In 1993, initial vegetation and conifer measurements and aspen removal within 1.8 m of the conifer was completed. These were followed by growth response measurements in 1994 and 1995. This progress report details the analysis of the second year lodgepole pine growth response.

Two years after treatment, there are significant and accelerating differences in radial growth response between treatments. The control plots consistently have the smallest radial growth; best growth is achieved under low levels of aspen competition with the $BDR > 0.75$ removal, followed closely by full removal. There are no significant differences in pine height growth between treatments. There is a trend toward higher mortality and mechanical and pest damage in plots where all the aspen have been removed, although at this point it is not statistically significant.

ACKNOWLEDGEMENTS

The project was designed with the assistance of Stan Navratil from the Canadian Forest Service, Edmonton. Roger Hayward from Weldwood of Canada, Hinton Division was involved in site selection, logistical and administrative support. Kent MacDonald of the Foothills Model Forest provided logistical and administrative support. Robert Lucas and his field assistants carried out the data collection in 1994 and 1995. Andrea Hagan assisted in data entry and production of this report. Accommodation was provided in 1995 by the Alberta Environmental Protection Land and Forest Service in Edson. The help of the above named individuals and organizations along with the summer students who provided data collection in 1993 is gratefully acknowledged.

Funding for this study was provided, in part, by Green Plan Canada, through the Foothills Model Forest.

TABLE OF CONTENTS

	Page
Abstract	ii
Acknowledgements	iii
Introduction	1
Study Location	2
Methods	2
Experimental Design	2
Plot Selection	4
Treatments	4
Field Measurements in 1993	5
Post-Treatment Field Measurements in 1994 and 1995	5
Analysis Methods	6
Results and Discussion	7
Vegetation Competition Levels Before and After Treatment in 1993	7
Aspen Competition Levels Two Years After Treatment	9
Pine Growth Response Two Years After Treatment	9
Mortality and Damage to the Target Lodgepole Pine	19
Conclusions	25
References	27
Appendix 1 Sample Data Collection Field Sheet for 1995	29

LIST OF FIGURES

1	Study Area and Experimental Layout of Blocks and Subblocks	3
2	Aspen Density Before and After Treatment - For All Blocks Combined	10
3	Aspen Height Before and After Treatment - For All Blocks Combined	11
4	Aspen Cover Before and After Treatment - For All Blocks Combined	12
5	Lodgepole Pine Root Collar Diameter Growth for All Blocks Combined - By Treatment	15
6	Lodgepole Pine Root Collar Diameter Growth for Block 1 - By Treatment	16
7	Lodgepole Pine Root Collar Diameter Growth for Block 2 - By Treatment	17
8	Lodgepole Pine Root Collar Diameter Growth for Block 3 - By Treatment	18
9	Lodgepole Pine Height Growth for All Blocks Combined - By Treatment	20
10	Lodgepole Pine Height Growth for Block 1 - By Treatment	21
11	Lodgepole Pine Height Growth for Block 2 - By Treatment	22
12	Lodgepole Pine Height Growth for Block 3 - By Treatment	23

LIST OF TABLES

1	Analysis of Covariance of Lodgepole Pine Growth in 1995, Using Pretreatment Size as a Covariant	13
2	Amount, Type and Severity of Damage to Target Lodgepole Pine Trees in 1995 for All Blocks Combined	24

INTRODUCTION

Performance expectations for juvenile conifers have been incorporated into the new free-to-grow regeneration standards in Alberta and extensive conifer release programs are implemented annually to bring regenerated stands to the provincially targeted standards. Selecting stands for the best response to and economic return from release treatments is difficult because of the high cost of treatment and limited information available on biological efficacy. Current treatment decisions are generally subjective or arbitrary and foresters require quantitative tools to assist in these decisions.

In 1993, the Canadian Forest Service completed a project on lodgepole pine-aspen competition. The objective of this study was to select or develop a competition index for quantifying the level of aspen competition that best predicts lodgepole pine growth. An index was required that would be easy to use in the field and applicable to release decisions. Based on this study, a new competition index, called the Basal Diameter Ratio (BDR) was developed (Navratil and MacIsaac 1993) which is a simplification of Lorimer's (1983) competition index:

$$CI = \frac{\text{tallest aspen basal diameter}}{\text{lodgepole pine basal diameter}}$$

Basal diameter refers to the stem diameter measured just above the root collar. In addition to its simplicity, it was as good or better in predicting pine response than other more complex competition indices, and has the potential to be used in an operational environment. Its potential has received favourable comment from operational foresters when presented at technical sessions, and they are eager to see that it is adopted.

This index was developed for lodgepole pine-aspen regeneration in west-central Alberta, but the study did not include actual release response assessment. The pine growth responses must be confirmed by field experiments. This Foothills Model Forest project was undertaken to ensure that this critical step is completed, before the index is used for stand tending decisions.

The purpose of this study is to test the application of the Basal Diameter Ratio (BDR) competition index developed by The Canadian Forest Service in tending decision to increase conifer growth. The goal of this study is to provide concrete data on how effective the BDR competition index is in guiding stand tending decisions in lodgepole pine-aspen blocks in west-central Alberta.

The study will help to answer the following questions:

1. How easy it is to apply the BDR competition index in determining what sections of a block should be targeted for stand tending?
2. At what level of aspen competition control (as quantified by the BDR index), is the best conifer growth achieved?

3. How consistent is the growth improvement in pine with a given level of removal of aspen competition?

A Foothills Model Forest report summarizing the project establishment and first year post-treatment results was produced by MacIsaac (1995). This progress report presents the second year post-treatment growth response of lodgepole pine following different levels of aspen removal based on the Basal Diameter Ratio competition index.

STUDY LOCATION

The study is located within the Weldwood Forest Management Area, within the Lower Foothills natural subregion¹. The Upper Foothills natural subregion was not suitable for this study because in that ecological zone, aspen is often not the major competitor of pine (willows, alder and balsam poplar are most dominant).

Three blocks were chosen for the study, based on field reconnaissance conducted in June and July 1993. These are in the Marlboro Working Circle, Compartment 8, Blocks 404, 378 and 378A, harvested in 1985 and 1986 (Fig 1). Specifically, these blocks met the following criteria; 1) Stand age between 8 and 13 years old (since clearcut). 2) At least 50% pine stocking. 3) At least 50% aspen stocking. 4) Not stand tended. 5) Planted within 3 years of harvest. The chosen blocks had excellent aspen and pine stocking with a minimal of other competitors. Initial field reconnaissance in 1993 indicated that the aspen were 2-3 m tall, and the pine were 0.5-1.0 m tall. All three blocks had been site prepared with a Bräcke scarifier prior to planting.

METHODS


Experimental Design

Four levels of aspen competition were established in 1993 by selectively removing aspen within 1.78 m of the pine trees (corresponding to a plot area of 10 m²), using the BDR index as a guide. The pine growth response was measured in 1994 and 1995, and will also be remeasured in 1996. A randomized nested design with three blocks is being used. Within each block, three well distributed areas (subblocks) at least 1.0 ha each were sampled (Fig. 1). In each subblock, 40 lodgepole pine-centred plots (1.78m radius) were established. The four treatments (listed below) were randomly assigned to the 40 plots; 10 plots received each treatment in each subblock. Once 10 plots of one treatment type were chosen, no more of that type were used in the subblock. A randomized complete

¹ Alberta natural regions from: Alberta. 1994. Natural regions and subregions of Alberta. [natural region map]. Land Information Services Division, Alberta Environmental Protection, Edmonton, Alberta. 1 sheet.

Figure 1

Study Area and Experimental Layout of Blocks and Subblocks



block design, in which each subblock would receive a set treatment for all plots within it was not used, because site conditions within each block were not uniform (a requirement for randomized complete block design). This randomized nested method allowed the interspersal of all four treatments throughout the blocks, thus removing bias.

Plot Selection

Placement of the permanent competition plots was based on a uniform grid within each subblock with a random starting point. Plots were spaced on a 10m by 10m grid. At each point on the grid, the closest lodgepole pine tree was used for the plot centre. If a suitable pine tree with aspen competition was not found within 5 m of the sampling grid, then there was no plot placed at that grid point. The plots had a fixed 1.78 m radius (10m²). Selected saplings were replaced with the next closest pine tree if damage unrelated to competition pressure was noted. A target tree² was selected if it had no recent moderate to severe damage due to herbivory, the leader growth was undamaged; had at least 5 internodes (to ensure that recent arrivals not used); was not advanced regeneration; had no major insect damage; and was not subject to significant intraspecific (pine-pine) competition (i.e. no crown overlap with other pines).

There were additional selection criteria, to ensure there was sufficient aspen competition. There was at least one aspen competitor, in at least three of the four quadrants around the pine tree, with a basal diameter equal to or larger than the pine tree (BD Ratio greater than 1). An aspen density criterion was also used, with a minimum of 8 aspen on the plot. Plots were not placed within 10 m of live residual trees. All target trees were selected using the above criteria.

Treatments

Each plot received one of the four treatments listed below.

- a) no aspen removal (control plot)
- b) removal of all aspen within 1.8 m of the target lodgepole pine tree where the aspen basal diameter was larger than the pine (BD ratio between pine and tallest remaining aspen was less than 1.0).
- c) removal of all aspen within 1.8 m of the target lodgepole pine tree where the aspen basal diameter was 75% of the pine diameter or greater (BD ratio between pine and tallest remaining aspen was less than 0.75).
- d) removal of all aspen within 1.8 m of the target lodgepole pine tree, regardless of aspen size.

Aspen competition was removed using hand saws, following vegetation measurement in 1993. Aspen competition was allowed to regenerate in subsequent years, and will not be removed in 1996.

² In this report, the term "target tree" refers to the specific crop tree of interest in each plot centre.

Field Measurements in 1993

Within the 1.78 m radius plots, competition data was collected in August and September, 1993. This was mostly baseline data collection, made prior to aspen removal (measurements on aspen remaining after treatment were also made, as described below).

Individual tree measurements were collected for the following trees: target lodgepole pine, the tallest conifer in the plot, the conifer in the plot that was closest to the target tree, and the tallest and closest aspen in the plot prior to treatment. For the two partial aspen removal treatments, if there was a new closest and/or tallest aspen, they were also measured. Target tree measurements were: crown height, crown radius, total height, root collar diameter, estimated age based on the number of internodes, height increments for the previous five years (including the current year) and percent overtopping of the top one-third of the crown by competing vegetation. As well, any slight damage along with the causal mechanism was noted (trees were not selected for study if the damage was moderate to severe). Measurements for the nearest and tallest conifer and hardwood in the plot included: azimuth from target tree, crown radius, total height, root collar diameter, distance from the target stem-to-inside crown of competitor, distance from the target stem-to-stem of competitor, distance from target stem-to-outside crown of competitor, estimated age based on internode counts and height increments for the previous five and three years, for conifers and hardwoods, respectively (including the current year). Descriptions of the detailed tree measurement variables are in MacIsaac (1995).

Aspen competition within each plot was measured both before and after the treatments in 1993. This included: average height and cover, total density and density by quadrant. Average height and density was also measured for each tree species, all trees combined, shrubs, forbs and grass. Microsite conditions within the plots were measured for the following variables: moisture class, drainage class, microtopography, litter depth, aspect, slope, slope position, and slash abundance.

Post-Treatment Field Measurements in 1994 and 1995

Within the 1.78 m radius plots, the pine growth response to the treatments and hardwood competition data was collected in August of 1994 and 1995, after the pine growth had stopped (lammas shoot growth, which is a second period of shoot elongation late in the summer under favourable moisture conditions, was not noted in the study blocks in those two years). Detailed measurements on the target tree and closest conifer and hardwood in the plot were identical as for 1993, except that crown radius was only measured on the target tree. The measurement of closest and tallest conifer and hardwood in the plots was done independently each year. In other words, due to differential growth and mortality of individuals, a tree selected as the closest and/or tallest in one year, might not be the tallest or closest the next year.

Average cover, height and density of hardwood trees was collected in the plots. Aspen density was recorded in two ways: 1. counting all individual shoots 2. lumping any shoots coppiced from a single aspen stem cut in 1993 as one shoot. Aspen density was collected for the whole plot, and for

each quadrant. In addition, a few other variables related to the target tree were collected: etiolation, herb crowding, vigour, occurrence and severity of disease, insect or mechanical damage. Density and average height and cover of each tree species in the plot was recorded, but unlike 1993, data on other growth forms and on microsite conditions was not recorded.

An example of the 1995 data sheet is in Appendix 1. Descriptions of the detailed tree measurement variables are the same as for 1993 and are described in MacIsaac (1995).

Analysis Methods

There were three types of analysis used, as follows:

- a) General statistical summaries and tests for normality and data transformations.
- b) Analysis of covariance to test differences in pine growth response two years after treatment.
- c) Multiple means tests to test for pine growth response differences and hardwood competition level differences two years after treatment.

This analysis was performed using the SAS statistical software package (SAS Institute Inc. 1989).

A variety of data transformations were used in an attempt to normalize the data prior to analysis, following the approach outlined in Sabin and Stafford (1990) and Zar (1984). The following transformations were tested: square root, square root of value+0.5, inverse of value+1, and natural log of value+1. The W-test for normality (Shapiro and Wilk 1965) as extended by Royston (1982) for sample sizes less than 2000 was used for all the variables. Tests for normality were performed on subpopulations based on stratification by treatment (n=90), and treatment by block (n=30). For all variables, there were specific transformations which consistently improved the distribution towards normality. Based on the above, the following transformations were used in the analyses: a) pine height increment and radial increment: no transformation b) pine height and root collar diameter: natural logarithm of value+1

For analysis of growth variables after treatment, covariance analysis was used, which included the size of the conifer prior to treatment (i.e., at the end of the 1993 growing season) (Woollons and White 1988). Cumulative two-year growth response was analyzed rather than treating each year's growth independently because the cumulative growth response was the primary variable of interest. Examining total growth response is a more elegant approach. It results in a more robust model, which is not affected by problems associated with independence of sequential measurements when yearly growth is examined separately. A mixed (fixed and random effects) linear model was developed, as appropriate for the experimental design (Borders and Shiver 1989; Neter et al. 1989; SAS Institute Inc. 1991). The model was:

Growth = tree size prior to treatment+block+subblock(block)+removal+removal*block+error

In this model, subblock was considered to be a random effect. Complete model statistics are presented in this report, following the recommendation of Warren (1986).

Ryan's multiple range test (c.f., Day and Quinn 1989) was used to determine significant differences in two-year post-treatment 1994-95 pine growth response and differences in the hardwood competition levels each year before and after treatment. These tests are preferable to the more commonly used Duncan's multiple range tests (Chew 1976; Jones 1984; Mize and Schultz 1985), for this experimental situation.

In 1993, during block selection and plot layout, some minor damage to regenerating lodgepole pine trees in the area was noted (MacIsaac 1995). Target trees were then selected which had no damage or only minor damage. There was a concern, however, that the damage might become more severe over time. Based on this consideration, a subjective classification of damage severity (none, slight, moderate, severe) and the damaging agent or physical sign of damage was recorded for each target tree. Damage to trees was classified into two groups: 1) Causes and signs of damage that were known or inferred to be related to insect and disease. This included root rot (*Armillaria* sp.), needle cast, chlorosis, western gall rust (*Endocronartium harknessii*), pitch blister moth *Petrova albicapitana*, stalactiform blister rust (*Cronartium* sp.) and resinosis. 2) Damage that was known or inferred to be related to mechanical damage or browse. This included: broken/damaged leader, broken/damaged branches, damaged base/stem (girdling), damaged roots, direct evidence of browsing, double top, damaged stem or forked stem.

Non-parametric analysis was done to test whether there was a relationship between target tree damage and removal of aspen (treatment). The Kruskal-Wallis rank-sum test was used to determine if there was a difference in the target tree damage between treatments, based on an average level of damage per tree in each plot. The following rank-sum tests were performed to determine the following:

1. If the aspen removal (treatment) had an effect on the level of occurrence (for any severity level) of insect and disease damage for target pine trees.
2. If the aspen removal (treatment) had an effect on the level of occurrence (for any severity level) of physical damage for target pine trees.
3. If the aspen removal (treatment) had an effect on the severity of damage (insect, disease and physical damage combined) for target pine trees.
4. If the aspen removal (treatment) had an effect on the mortality rate (all causes combined) for target pine trees.

RESULTS AND DISCUSSION

Vegetation Competition Levels Before and After Treatment in 1993

The regenerating blocks chosen for this study had been planted to lodgepole pine. Aspen was the major competitor, with much lower amounts of shrub, forb and grass competition. Prior to treatment, the 2-3 m tall aspen was moderately-dense (ranging from 21 to 32 thousand stems/ ha), with an average cover of 38-50%. The planted lodgepole pine was one-third to one-half the aspen height (82-

138 cm). The grass competition, which was primarily bluejoint (*Calamagrostis canadensis*) with lesser amounts of wild hairy rye (*Elymus innovatus*), was moderate to light, with cover ranging from 9 to 16%. The shrub strata was low, with the majority of cover below 50 cm. In terms of cover, density and height, balsam fir was the second most dominant conifer species, after pine. As well, the blocks had minor amounts of white spruce and paper birch.

Analysis of variance of aspen density, height and cover for blocks and subblocks indicated that the aspen competition was not uniform throughout the blocks prior to treatment. For density and height, aspen had more within-block variability than between-block variability. Least squares means tests showed significantly greater density (31,800 stems/ha) in block 3 compared to the other two blocks ($P < 0.05$). This was due to a high density in block 3, subblock 3 of 43,400 stems/ha, which was almost double the density for most of the other subblocks. For aspen cover, block 2 had significantly less cover than blocks 1 and 3, but the differences were not as large as for aspen density. Aspen height differences between blocks was less pronounced than for the other two aspen variables, although block 1 did have significantly taller aspen (263 cm) than blocks 2 and 3. Overall, in terms of aspen competition, the only "outlier" was for aspen density in block 3, subblock 3. These analyzed differences in aspen competition within blocks supported the decision to mix all four treatments within each of the subblocks. When all blocks were combined, there were no significant differences in aspen density before treatment, averaging around 25,000 stems/ha.

When mean values were generated based on treatment type (with plots of each treatment evenly distributed throughout each subblock), there were only small differences in pre-treatment aspen competition levels. Removal of aspen based on the BDR of 1.0 (i.e., all aspen with a root collar diameter greater than that of pine), resulted in a 47-66% decrease within blocks in aspen density, and 50-66% decrease in average aspen height within blocks. For aspen cover, the post-treatment decrease within blocks was more pronounced (87-92%), because the removal took out the larger trees. Using a more stringent BDR of 0.75 as a guide did not result in an appreciably greater removal of aspen. This is because, on average, the dominant aspen competitors would have a greater root collar diameter than the pine, due to faster growth rates of aspen as compared to pine in regenerating stands (and so for a given plot, using a BDR of 1.0 would achieve similar post-treatment levels of aspen as would thinning based on a BDR of 0.75). Using a more stringent removal criteria of 0.75 would result in a large number of smaller stems to be removed, which would not influence the average remaining competition very much.

An analysis of variance was performed on the average aspen competition after removal, to quantify the amount of variation within and between treatments and blocks. As expected, removal (treatment) had a very significant effect on aspen density, height and cover (Figs. 2 to 4). This is mostly due to the effect of the two extreme treatments (all removal versus no removal). Least squares means tests indicated, that, in most blocks, there were no significant difference in the remaining level of aspen for the two intermediate treatments.

Aspen Competition Levels Two Years After Treatment

Figures 2-4 provide a graphical summary of aspen density, average aspen height and average aspen cover before and after treatment. Aspen density counts which include coppiced stems showed significant regeneration in the first year after removal, with densities approaching the pre-harvest levels, especially for the plots with all aspen removed (Figs.2). However, by the second year, densities in all three treatments had fallen significantly below that of the control plots. In contrast to density, average aspen height and cover in 1994 and 1995 remained well below the levels recorded in 1993 before treatments (Figs. 3 and 4).


Pine Growth Response Two Years After Treatment

The post-treatment analysis is presented using three complementary techniques: 1. covariance analysis, 2. Graphical representation of the growth trajectories, and 3. multiple means tests. Because of the effect of the conifer tree size prior to treatment, analysis of covariance was required when analyzing the second year growth response to treatment. The results of analysis of covariance, which controlled for differences in the initial size of the pine, are presented in Table 1. As expected, the covariant had a significant effect on all growth variables ($P < 0.0001$). For 1995 height and cumulative 1994-95 height increment, there were no treatment effects. This is confirmed with the least squares means tests which indicates no significant differences between any of the treatment responses, except for Block 2 for 1995 Height. In this latter case, the intermediate treatment where plots with aspen greater than a BDR of 0.75 was removed had significantly larger height growth.

For root collar diameter and radial increment, there were significant removal effects ($P < 0.0001$). The least squares means tests indicate that the significant differences were usually due to lower growth response for the control (no removal) compared to the other treatments. There were no differences in growth response from the intermediate removals based on BDR values of 1.0 and 0.75. This is corroborated by the results of Ryan's Multiple means tests (results not shown). For increment in root collar diameter, in all three blocks, the average rate of radial growth was least for the treatment where no aspen was removed.

Figures 5 to 8 present a graphical perspective of the two year post-treatment growth response for pine root collar diameter. The trends which were evident in the first year post-treatment response (MacIsaac 1995) have become more significant after the second year of response. Consistently, for all blocks combined, and each block separately, the smallest root collar diameter growth was in the control plots with no aspen removal. For all blocks combined and in block 2, this difference was significant. The best root collar diameter growth is in plots where there is a low level of aspen retained (aspen with BDR > 0.75 removed), although the differences with the other treatments are not significant. The level of aspen associated with this treatment is quite modest: two years after treatment, the average aspen height was only half that of the pine (99 cm vs 184 cm), with an associated aspen cover of only 7%. This can hardly be considered high aspen competition. The critical factor is whether the remaining aspen will eventually overtop the pine.

Figure 2
Aspen Density Before and After Treatment - For All Blocks Combined





Analysis based on all blocks combined (n= approx. 90 1.78 m radius plots for each treatment). Standard error of the mean shown (upper limit only).

Similar letters for the same year indicate means not significantly different, at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc. 1990).

Figure 3

Aspen Height Before and After Treatment - For All Blocks Combined





Analysis based on all blocks combined (n=approx. 90 1.78 m radius plots for each treatment). Standard error of the mean shown (upper limit only).

Similar letters for the same year indicate means not significantly different, at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc. 1990).

Figure 4

Aspen Cover Before and After Treatment - All Blocks Combined



Analysis based on all blocks combined (n=approx. 90 1.78 m radius plots for each treatment). Standard error of the mean shown (upper limit only).

Similar letters for the same year indicate means not significantly different at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc. 1990).

Table 1
 Analysis of Covariance of Lodgepole Pine Growth in 1995
 Using Pretreatment Size as a Covariant

1995 Height¹				
Source	DF	Mean Sq.	F Value	Pr > F
1993 Height ²	1	25.38	1227	0.0001
Block ²	2	0.05	2.1316	0.1822
Subblock (Block) ³	6	0.03	1.6174	0.1415
Removal	3	0.02	1.2346	0.2971
Block * Removal	6	0.03	1.5826	0.1514
Error	330	0.02		
Least Squares Means (Values are means (cm) ± standard error of the mean)				
Removal	All	0.75	1.00	None
Block 1	211.41 ±7.89a	220.80 ±11.30a	224.76 ±11.63a	212.48 ±10.07a
Block 2	157.67 4.71ab	168.00 7.45a	140.50 6.34b	153.31 7.96ab
Block 3	147.10 7.11a	160.59 6.10a	148.97 7.07a	158.41 6.85a
All Blocks Combined	171.62 4.84a	183.74 5.74a	171.06 6.38a	173.85 5.55a

Cumulative 1994-1995 Height Increment				
Source	DF	Mean Sq.	F Value	Pr > F
1993 Height	1	33321.94	46.4532	0.0001
Block	2	2313.65	2.3147	0.1685
Subblock (Block)	6	1135.31	2.7214	0.0136
Removal	3	687.58	1.6481	0.1781
Block * Removal	6	419.12	1.0046	0.4221
Error	330	417.19	-	-
Least Squares Means (Values are means (cm) ± standard error of the mean)				
Removal	All	0.75	1.00	None
Block 1	69.17 ±4.91a	72.97 ±4.35a	74.13 ±5.05a	69.04 ±5.12a
Block 2	64.90 3.11a	70.35 4.32a	59.50 4.01a	65.31 4.44a
Block 3	54.43 4.07a	64.44 3.99a	64.10 3.89a	68.48 3.39a
All Blocks Combined	62.76 2.42a	69.29 2.44a	65.84 2.56a	67.58 2.49a

Table 1 (continued)

1995 Root Collar Diameter

Source	DF	Mean Sq.	F Value	Pr > F
1993 RCD	1	22.65	1010	0.0001
Block	2	0.05	2.2197	0.1783
Subblock (Block)	6	0.02	1.0763	0.3763
Removal	3	0.27	12.2530	0.0001
Block * Removal	6	0.01	0.5177	0.7949
Error	328	0.02	-	-

Least Squares Means (Values are means (mm) ± standard error of the mean)				
Removal	All	0.75	1.00	None
Block 1	34.97 ±1.59b	36.53 ±2.08b	36.28 ±1.84b	32.12 ±1.89a
Block 2	31.27 1.36b	30.82 1.49b	25.97 1.20b	26.14 1.32a
Block 3	26.10 1.46ab	27.48 1.03b	25.41 1.34b	24.18 1.15a
All Blocks Combined	30.73 0.92b	31.68 1.01b	29.18 1.00b	27.35 0.91a

Cumulative 1994-1995 Radial Increment

Source	DF	Mean Sq.	F Value	Pr > F
1993 RCD	1	187.95	35.3515	0.0001
Block	2	8.80	1.3522	0.3196
Subblock (Block)	6	6.66	1.3665	0.2274
Removal	3	68.53	14.0536	0.0001
Block * Removal	6	3.81	0.7815	0.5849
Error	340	4.88	-	-

Least Squares Means (Values are means (mm) ± standard error of the mean)				
Removal	All	0.75	1.00	None
Block 1	6.13 ±0.50b	6.13 ±0.43b	6.28 ±0.51b	4.29 ±0.38a
Block 2	5.70 0.46b	6.16 0.38b	5.33 0.38b	3.83 0.33a
Block 3	4.67 0.41a	4.93 0.30a	4.69 0.31a	3.79 0.31a
All Blocks Combined	5.49 0.27b	5.74 0.22b	5.43 0.24b	3.96 0.20a


¹Analysis performed with natural log (value+1) transformation for height and RCD; no transformation for height increment and radial increment.

²Tests of hypotheses use the Type I MS for Subblock(Block) as an error term.

³Subblock is designated as a random effect.


⁴Means in each row followed by the same letter do not differ significantly (P≥0.05) in least squares means test.

Figure 5
Lodgepole Pine Root Collar Diameter Growth for All Blocks Combined
By Treatment



Standard error of the mean shown (lower interval only).
 Similar letters for the same year indicate means not significantly different,
 at P=0.05 using Ryan et al's Multiple Range Test.

Figure 6
Lodgepole Pine Root Collar Diameter Growth for Block 1 - By Treatment




- All aspen removed
 - Aspen with BDR > 0.75 removed
 - ▲—▲ Aspen with RCD > 1.0 removed
 - ▼—▼ No aspen removed
- n = approx. 30 for each treatment

Standard error of the mean shown (lower interval only).

Means not significantly different for removal type in all years shown, at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc.1990).


Figure 7
Lodgepole Pine Root Collar Diameter Growth for Block 2 - By Treatment



Standard error of the mean shown (lower interval only).

Similar letters for the same year indicate means not significantly different, at P=0.05 using Ryan et al's Multiple Range Test.

Figure 8
Lodgepole Pine Root Collar Diameter Growth for Block 3 - By Treatment



- All aspen removed
 - Aspen with BDR > 0.75 removed
 - ▲—▲ Aspen with RCD > 1.0 removed
 - ▼—▼ No aspen removed
- n = approx. 30 for each treatment

Standard error of the mean shown (lower interval only).
Means not significantly different for removal type in all years shown,
at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc.1990).

Height growth has not responded to the different treatments. In general, the relative performance after two years is mostly a function of the differences in average size prior to treatments (Figs.9-12). The growth curves have not diverged and are not significantly different after two years. These results clearly indicate the effect of initial pine size on the short term average growth response of pine height growing under post-treatment conditions.

The results for radial growth clearly indicate that the divergent growth trajectories that were initiated in the first year after treatment have become more pronounced after the second year. For example, Figure 5 and 6 show there is divergence in average root collar diameter for trees growing under no aspen removal, compared to the other treatments. The fact that radial increment showed an increasing treatment response after two years year while height increment has not is consistent with findings from other studies on aspen-pine competition. Navratil and MacIsaac (1993) indicate that released pine respond with increased radial growth prior to any observed increase in height growth. Juvenile spacing trials in the region show that spacing improves diameter growth on all sites types, but that height growth is enhanced only on poor to medium sites (Johnstone 1981, Yang 1991). These Marlboro sites are similar to other locations in the region that would be considered to be of good site quality (Duffy 1964). As the pines respond to release, carbon allocation would be directed to roots and radial growth before height growth.


It is also intuitive that the differences in growth response would be significant when comparing the controls against the aspen removal treatments, as the three aspen removal treatments have resulted in aspen abundances that are much more alike then the controls. It will probably take more time for growth responses to become noticeable with the intermediate treatments. The lack of differences in growth response between the two intermediate treatments may also be because, in some cases, there was not a significant difference in the remaining aspen competition between the two intermediate treatments.

Mortality and Damage to the Target Lodgepole Pine

The amount, severity and type of damage to target lodgepole pine trees is summarized in Table 2. Sixty-five percent of all target trees in the three blocks combined did not show any signs of damage or disease. In damaged trees, *Armillaria* root rot fungus was noted as one of the major causes of damage (8.6% of the total target trees). Between the 1993 and 1994 field seasons, 11 target trees died. From 1994 to 1995 an additional 25 target trees died, for a cumulative mortality of 10%. Most of the mortality and severe damage was due to *Armillaria* root rot, with lesser amounts due to western gall rust and (Table 2). This mortality rate is approximately five times greater than the average for lodgepole pine growing on medium to high productivity sites in the area (Ives and Rentz 1993).

There is a concern that this mortality could increase substantially over time. While the major *Armillaria* food source is probably stumps from the preharvested stand, the aspen cut in 1993 could exacerbate this trend, through the inclusion of dead aspen stumps and stems as an additional food

Figure 9
Lodgepole Pine Height Growth for All Blocks Combined - By Treatment




- All aspen removed
- Aspen with BDR > 0.75 removed
- ▲—▲ Aspen with RCD > 1.0 removed
- ▼—▼ No aspen removed

n = approx. 90 for each treatment

Standard error of the mean shown (lower interval only).

Means not significantly different for removal type in all years shown, at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc.1990).

Figure 10
Lodgepole Pine Height Growth for Block 1 - By Treatment




- All aspen removed
- Aspen with BDR > 0.75 removed
- ▲—▲ Aspen with RCD > 1.0 removed
- ▼—▼ No aspen removed

n = approx. 30 for each treatment

Standard error of the mean shown (lower interval only).
Means not significantly different for removal type in all years shown.
at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc.1990).


Figure 11
Lodgepole Pine Height Growth for Block 2 - By Treatment



Standard error of the mean shown (lower interval only).

Similar letters for the same year indicate means not significantly different, at P=0.05 using Ryan et al's Multiple Range Test.

Figure 12
Lodgepole Pine Height Growth for Block 3 - By Treatment



- All aspen removed
- Aspen with BDR > 0.75 removed
- ▲—▲ Aspen with RCD > 1.0 removed
- ▼—▼ No aspen removed

n = approx. 30 for each treatment

Standard error of the mean shown (lower interval only).
Means not significantly different for removal type in all years shown.
at P=0.05 using Ryan et al's Multiple Range Test (SAS Institute Inc. 1990).

Table 2
Amount, Type and Severity of Damage to Target Lodgepole Pine Trees
in 1995 for All Bloks Combined

Most Prevalent Damage ³	Damage Severity ^{1,2}				Total		Dead Trees
	None	Slight	Moderate	Severe	N	%	
No Damage	235	.	.	.	235	65.3	1
Root Rot (<i>Armillaria</i> sp.)	.	.	.	31	31	8.6	31
Broken/Damaged Leader	.	6	1	1	8	2.2	.
Broken/Damaged Branches	.	3	.	.	3	0.8	.
Needle Cast (Undifferentiated)	.	2	2	.	4	1.1	.
Damaged Base/Stem (Girdling)	.	.	1	.	1	0.3	.
Chlorosis (Cause not defined)	.	7	13	7	27	7.5	.
Roots	.	.	.	1	1	0.3	1
Western Gall Rust (<i>Endocronartium harknessii</i>)	.	2	.	4	6	1.7	.
Double Top	.	1	.	.	1	0.3	.
<i>Petrova albicapitana</i>	.	1	1	1	3	0.8	.
Stalactiform Blister Rust (<i>Cronartium</i> sp.)	.	.	.	3	3	0.8	1
Browsed	.	20	7	1	28	7.8	.
Stem	.	3	1	.	4	1.1	.
Forked	.	1	.	.	1	0.3	.
Resinosis	.	1	.	.	1	0.3	.
Unknown (Dead)	.	.	.	3	3	0.8	2
Total	235	47	26	52	360	100	36

1. Based on a subjective evaluation.
2. Intermediate categories were moved up to the higher category.
3. While trees may have had damage from multiple sources, only the dominant damage was recorded.
4. A "." indicates none in that category.

source for the fungus (Dr. Ken Mallet, pers. comm). Over the three year period of the study the additional dead aspen material may not substantially influence the spread of *Armillaria*.

Other causes or signs of damage were (in decreasing importance): browse damage, chlorosis (undefined cause), physical damage to the leader and western gall rust. The major diseases found in these blocks has been reported for similar regenerating pine stands in the region (Bella 1985a, Ives and Rentz 1993).

The four Kruskal-Wallis rank-sum tests all indicated that the amount of damage or mortality of target trees was approximately double (though still not widespread) for the plots where all the aspen was removed, compared to the control plots. However, there was a lot of variation in this relationship, as is shown by the non-significant probability level associated with each rank-sum test:

1. H_0 : Aspen removal (treatment) had an effect on the level of occurrence (for any severity level) of insect and disease damage for target pine trees (not significant: $P=0.09$).
2. H_0 : Aspen removal (treatment) had an effect on the level of occurrence (for any severity level) of physical damage for target pine trees (not significant: $P=0.36$).
3. H_0 : Aspen removal (treatment) had an effect on the severity of damage (insect, disease and physical damage combined) for target pine trees (not significant: $P=0.06$).
4. H_0 : Aspen removal (treatment) had an effect on the mortality rate (all causes combined) for target pine trees (not significant: $P=0.68$).

Thinned stands may lead to increased incidence of pest damage (e.g., Bella 1985a, 1985b), possibly through such mechanisms as increased wind dispersal of spores. It is possible that removal of all aspen adjacent to the pine may exacerbate similar pest problems. The factors that influence damage to pine may act in different ways. For instance, Bella (1985a) and Ives and Rentz (1993) indicate that mammal damage in young lodgepole pine is greater in more dense stands than in less dense ones, however, pest damage may be greater in thinned stands (Bella 1985a). There is some early evidence from this study that retention of low levels of aspen may be associated with optimal growth of pine, in cases where the aspen is shorter than the pine. As the aspen crowns develop, second removal may be required to ensure optimal pine growth.

CONCLUSIONS

1. The aspen competition levels prior to treatment in 1993 were uniform, except for subblock 3 in block 3 which had a significantly greater density of aspen.
2. The second year of post-treatment pine growth showed a continuation of trends seen one year after treatment. After two years of post-treatment growth, radial increment and total root collar diameter is significantly lower in the control plots. The best radial growth is associated with plots under low levels of aspen competition with aspen with a BDR of more than 0.75 removal, followed closely by full removal.

3. After two years of post-treatment growth, there were no significant differences in height growth between any of the treatments.
4. There is a trend toward higher mortality and mechanical and pest damage in plots where all the aspen have been removed, although two years after treatment it is not statistically significant.

REFERENCES

- Bella, I.E. 1985a. Pest damage incidence in natural and thinned lodgepole pine in Alberta. *For. Chron.* 61:233-238.
- Bella, I.E. 1985b. Western gall rust and insect leader damage in relation to tree size in young lodgepole pine in Alberta. *Can. J. For. Res.* 15:1008-1010.
- Borders, B.E.; Shiver, B.D. 1989. Herbicide field studies in forestry: Statistical and other considerations. *Can. J. For. Res.* 19:768-772.
- Chew, V. 1976. Comparing treatment means: A compendium. *Hortscience.* 11(4):348-356.
- Day, R.W.; Quinn, G.P. 1989. Comparisons of treatments after an analysis of variance in ecology. *Ecological Monographs.* 59(4):433-463.
- Duffy, P.J.B. 1964. Relationship between site factors and growth of lodgepole pine (*Pinus contorta* Dougl. var. *laifolia* Engelm.) in the foothills section of Alberta. Dept. Forestry., Publication No. 1065.
- Ives, W.G.H; Rentz, C.L. 1993. Factors affecting the survival of immature lodgepole pine in the foothills of west-central Alberta. *For. Can., Northwest Reg., North. For. Cent., Edmonton, Alberta. Info. Rept. NOR-X-330.*
- Johnstone, W.D., 1981. Effects of spacing 7-year-old lodgepole pine in west-central Alberta. *Can. For. Serv., Northwest Reg., North. For. Cent., Edmonton, Alberta. Info. Rept. NOR-X-236.*
- Jones, D. 1984. Use, misuse, and role of multiple-comparison procedures in ecological and agricultural entomology. *Forum: Environmental Entomology.* 13(3):635-648.
- Lorimer, C.G. 1983. Tests of age-independent competition indices for individual trees in natural hardwood stands. *For. Ecol. Manage.* 6:343-360.
- MacIsaac, D.A. 1995. Validation of basal diameter ratio competition index for lodgepole pine-aspen. Establishment and progress report. Foothills Model Forest. Hinton, Alberta.
- Mize, C.W.; Schultz, R.C. 1985. Comparing treatment means correctly and appropriately. *Can. J. For. Res.* 15:1142-1148.
- Navratil, S.; MacIsaac, D.A. 1993. Competition index for juvenile mixed stands of lodgepole pine and aspen in west-central Alberta. *For. Can., Northwest Reg., North. For. Cent., Edmonton, Alberta. For. Manage. Note 57.*

- Neter, J.; Wasserman, W.; Kutner, M.H. 1989. Applied linear regression models. Irwin, Boston, Massach. U.S.
- Royston, J.P. 1982. An extension of Shapiro and Wilk's W test for normality to large samples. Applied Statistics 31:115-124.
- Sabin, T.E.; Stafford, S.G. 1990. Assessing the need for transformation of response variables. Forest Research Laboratory, Oregon State University, Corvallis. Special Publication 20. 31 p.
- SAS Institute Inc. SAS/STAT User's Guide, version 6, fourth edition, volume 2. Cary, NC: SAS Institute Inc. 1989. 846 p.
- SAS Institute Inc. SAS System for Linear Models. Third edition. Cary, NC: SAS Institute Inc. 1991. 329 p.
- Shapiro, S.S.; Wilk, M.B. 1965. An analysis of variance test for normality (complete samples). Biometrika 52:591-611.
- Shaw, R.G.; Mitchell-Olds, T. 1993. Anova for unbalanced data: An overview. Ecology, 74(6);1638-1645.
- Warren, W.G. 1986. On the presentation of statistical analysis: reason or ritual. Can. J. For. Res. 16:1185-1191.
- Woollons, R.C.; White, A.G.D. 1988. Multiple covariance: Its utility in analysing forest fertilizer experiments. Forest Ecology and Managements. 25:59-72.
- Yang, R.C.. 1991. Early stand development of lodgepole pine spaced at age 7 in west-central Alberta. For. Can., Northwest Reg., North. For. Cent., Edmonton, Alberta. Info. Rept. NOR-X-322.
- Zar, J.H. 1984. Biostatistical analysis. Prentice-Hall, Englewood Cliffs, New Jersey.

Appendix 1
 Sample Data Collection Field Sheet for 1995

1995 BASAL DIAMETER RATIO CI VALIDATION (FOOTHILLS FOREST)

Block __ Subblock __ Plot __ Date: ____ 1995 Observer: _____ Recorder: _____

Aspen Removal: ____ (1=All 2=BDR >1.0 3=BDR >0.75 4=None)

Etiolation ____ Herb/Shrub Crowding ____ Subject Tree Vigour: ____

Disease/Insect/Mechanical Damage:Severity: ____ Type/Where: _____

PLOT INFORMATION

<p style="text-align: center;">Trees (all sizes)</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 25%;">Species Num</td> <td style="width: 10%;">Ht</td> <td style="width: 10%;">Cov</td> <td style="width: 25%;">Species Num</td> <td style="width: 10%;">Ht</td> <td style="width: 10%;">Cov</td> </tr> <tr> <td>P1</td> <td>_____</td> <td>_____</td> <td>Pb</td> <td>_____</td> <td>_____</td> </tr> <tr> <td>P12</td> <td>_____</td> <td>_____</td> <td>Fb</td> <td>_____</td> <td>_____</td> </tr> <tr> <td>At</td> <td>_____</td> <td>_____</td> <td>Sw</td> <td>_____</td> <td>_____</td> </tr> <tr> <td>_____</td> <td>_____</td> <td>_____</td> <td>All</td> <td>_____</td> <td>_____</td> </tr> </table> <p>P1 includes subject tree. P12 and All exclude subject tree</p>	Species Num	Ht	Cov	Species Num	Ht	Cov	P1	_____	_____	Pb	_____	_____	P12	_____	_____	Fb	_____	_____	At	_____	_____	Sw	_____	_____	_____	_____	_____	All	_____	_____	<p style="text-align: center;">Aspen Density in Quadrants</p> <table style="width: 100%; text-align: center;"> <tr> <td>N</td> <td>E</td> <td>S</td> <td>W</td> </tr> <tr> <td>_____</td> <td>_____</td> <td>_____</td> <td>_____</td> </tr> </table> <p>Count stems on stump as 1, then count as individuals (in brackets).</p>	N	E	S	W	_____	_____	_____	_____
Species Num	Ht	Cov	Species Num	Ht	Cov																																		
P1	_____	_____	Pb	_____	_____																																		
P12	_____	_____	Fb	_____	_____																																		
At	_____	_____	Sw	_____	_____																																		
_____	_____	_____	All	_____	_____																																		
N	E	S	W																																				
_____	_____	_____	_____																																				

SUBJECT, TALLEST AND CLOSEST CONIFER AND HARDWOOD (ALWAYS INCLUDE ASPEN)

	Sp	Cd	Az	Ht (cm)	Crown Ht Rad (cm)	% OT	RCD (mm)	STIN Dist (cm)	STST Dist (cm)	STOUT Dist (cm)	Height Increments (cm)				
											95	94	93	92	91
Con	P1	1		_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
	—	—	—	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
	—	—	—	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
Hdwd	—	—	—	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
	—	—	—	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____
	—	—	—	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____	_____