

Coarse Woody Debris in the Coastal Forests of Southern Vancouver Island

Ralph W. Wells, Centre for Applied Conservation Biology, University of British Columbia, 270-2357 Main Mall, Vancouver, British Columbia, Canada, V6T 1Z4

and

J.A. Trofymow, Pacific Forestry Centre, Canadian Forest Service, Natural Resources Canada, 506 West Burnside Road, Victoria, British Columbia, Canada, V8Z 1M5

Pages 23-24 *in* J.A. Trofymow and A. MacKinnon, editors. Proceedings of a workshop on Structure, Process, and Diversity in Successional Forests of Coastal British Columbia, February 17-19, 1998, Victoria, British Columbia. Northwest Science, Vol. 72 (special issue No. 2).

Acrobat version prepared and distributed by:

Natural Resources Canada
Canadian Forest Service
Pacific Forestry Centre
506 West Burnside Road
Victoria, British Columbia
V8Z 1M5
Canada

Reprinted with permission from Northwest Science, Volume 72, Special Issue No. 2, Washington State University Press, 1998.



Natural Resources
Canada

Canadian Forest
Service

Ressources naturelles
Canada

Service canadien
des forêts

Canada

and

J.A. Trofymow, Pacific Forestry Centre, Canadian Forest Service, Natural Resources Canada, 506 West Burnside Road, Victoria, British Columbia, Canada, V8Z 1M5

Coarse Woody Debris in the Coastal Forests of Southern Vancouver Island

In this extended abstract, we present results from a study of coarse woody debris in forests of coastal British Columbia (see Wells and Trofymow 1997 for the full report). We examined trends of CWD abundance across chronosequence plots established by the Canadian Forest Service, on the drier east side (CWHxm subzone; Meidinger and Pojar 1991) and the wetter west side (CWHvm subzone) of

southern Vancouver Island. Overall, total biomass of CWD on the east side was substantially (and significantly) lower than on the west side. Mean total biomass, across the chronosequences, ranged from 17 Mg/ha to 38 Mg/ha (55 m³/ha to 149 m³/ha) on the east side and 65 Mg/ha to 191 Mg/ha (307 m³/ha to 636 m³/ha) on the west side (Figures 1a and 1b).




Figure 1. Mean CWD mass (>12 cm dia.) for (a) east-side CWHxm and (b) west-side CWHvm chronosequences with decay class; mean CWD mass for (c) east-side CWHxm and (d) west-side CWHvm chronosequences with size class. Note change in scale between east-side and west-side plots. Error bars are standard errors. $n = 4$ plots/age class category. R = regeneration plots (3-9 years). I = immature plots (32-43 years). M = mature plots (66-99 years). O = old growth plots (>200 years). DC = decay class. SC = Size Class; SC 1 = 1.1 to 12.0 cm dia.; SC 2 = 12.1 to 29.9 cm dia.; SC 3 = 30.0 - 59.9 cm dia.; and SC 4 = 60 cm+ dia.

Spies and Franklin (1988) predicted that CWD would follow a 'U'-shaped trajectory in coastal forests of the Pacific Northwest. We found that total biomass of CWD followed a 'U'-shaped curve over the chronosequence in three of four sites on the west side, while no overall trend was detected for total biomass in the east side. Further, in both subzones, CWD was skewed to lower decay classes in younger stands, and more normally distributed in older stands (Figures 1a and 1b). On the east side, CWD in young stands tended towards small size classes, and was more uniformly distributed in older stands (Figure 1c; note that small diameter material is included). On the west side, CWD tended towards larger size classes in all age classes (Figure 1d; note that small diameter material is included). Finally, on the east side, Douglas-fir (*Pseudotsuga menziesii*) was the most common species among classified CWD; on the west side, western hemlock (*Tsuga heterophylla*) and western redcedar (*Thuja plicata*) were most common.

The biomass levels we observed on west side plots were consistent with accumulations reported in other *Pseudotsuga* - *Tsuga* forests of the Pacific Northwest (Caza 1993). Alternatively, on the drier east side plots, the range of 17 Mg/ha to 38 Mg/ha, (55 m³/ha to 149 m³/ha), were among the lowest CWD accumulations recorded for this forest type, regardless of age class. Reported biomass in old-growth (200+ years) *Pseudotsuga* - *Tsuga* forests of Washington and Oregon ranges from 66 Mg/ha to 490 Mg/ha (221 m³/ha to 1421 m³/ha) (Caza 1993).

The low amounts of CWD observed in this study on the east-side plots are somewhat surprising, given the high values observed in Douglas-fir forests in Washington and Oregon. It may be that the old-growth plots in this study were established in less productive stands than typical

for the east side (because productive sites in the CWHxm on southern Vancouver Island have been subject to more harvesting than less productive sites), yielding lower than typical CWD loadings. Decay rates, which can be high on dry sites (Harmon et al. 1986), combined with lower growth rates of trees could result in lower than typical CWD loadings on unproductive sites.

On both the east side and west side, variance in CWD loadings among plots was substantial. This may have limited our ability to detect some trends, and suggests that factors other than age (e.g. edaphic site characteristics and stochastic disturbance events) are influencing CWD abundance. Finally, we caution that the regenerating, immature, and most mature stands examined in this study may not be following the same developmental trajectory as old stands because they initiated after harvesting while old stands initiated after natural disturbances. Further studies examining the influence of site characteristics and disturbance history on CWD abundance would improve understanding of CWD dynamics in the CWH zone.

Acknowledgements

We thank Gordon Kayahara and an anonymous reviewer for helpful comments on this abstract. We also thank B. Blackwell and his field staff (B.A. Blackwell and Associates), B. Rowsell and Katherine Maxcy for their technical assistance. This work was supported in part by the ENFOR (ENergy From the FOREst) program of Forestry Canada, Projects P-404 and P-453; and by Forest Renewal BC, Award No. HQ96247 to J.A. Trofymow. Funding assistance by Forest Renewal BC does not imply endorsement of any statements or information contained herein.

Literature Cited

- Caza, C.L. 1993. Woody debris in the forests of British Columbia: a review of the literature and current research. BCMOF, Land Manage. Rep. 78. Victoria, BC.
- Harmon, M.E., J.F. Franklin, F.J. Swanson, P. Sollins, S.V. Gregory, J.D. Lattin, N.H. Anderson, S.P. Cline, N.G. Aumen, J.R. Sedell, G.W. Liendaemper, K. Cromack, JR., and K.W. Cummins. 1986. Ecology of coarse woody debris in temperate ecosystems. Adv. Ecol. Res. 15:133-303.
- Meidinger, D. and J. Pojar (eds.). 1991. Ecosystems of British Columbia. BCMOF, Research Branch Special Report, Series 6. Victoria, BC.
- Spies, T.A. and J.F. Franklin. 1988. Old growth and forest dynamics in the Douglas-fir region of western Oregon and Washington. Nat. Areas J. 8:190-201.
- Wells, R.W. and J.A. Trofymow. 1997. Coarse woody debris in chronosequences of forests on southern Vancouver Island. BC-X-375. Nat. Res. Can., Can. For. Ser., Pacific Forestry Centre, Victoria, BC.