

RESULTS OF FOREST INSECT AND  
DISEASE SURVEYS IN THE  
NORTHERN REGION OF ONTARIO  
1988

(FOREST DISTRICTS: MOOSONEE, HEARST, KAPUSKASING, COCHRANE  
CHAPLEAU, TIMMINS, KIRKLAND LAKE AND GOGAMA)

L.S. MacLEOD, V. JANSONS and S. PAYNE

FORESTRY CANADA  
ONTARIO REGION  
GREAT LAKES FORESTRY CENTRE  
1989

MISCELLANEOUS REPORT NO. 82

•Minister of Supply and Services Canada 1989  
Catalogue No. Fo29-8/82E  
ISBN 0-662-16709-0  
ISSN 0832-7130

Additional copies of this publication  
are available at no charge from:

Communications Services  
Great Lakes Forestry Centre  
Forestry Canada  
P.O. Box 490  
Sault Ste. Marie, Ontario  
P6A 5M7

Microfiches of this publication may be purchased from:

Micromedia Inc.  
Place du Portage  
165, Hôtel-de-Ville  
Hull, Quebec  
J8X 3X2

## **SURVEY HIGHLIGHTS**

This report documents the more important insect and disease problems that affected the forests of the Northern Region of Ontario during the 1988 field season, as determined by Forest Insect and Disease Survey (FIDS) personnel.

Several factors, including parasitism and disease, precipitated the collapse of the forest tent caterpillar outbreak in Kirkland Lake District but a new outbreak appears imminent in Hearst District. The birch leafminer continued to cause severe foliar damage in white birch stands in most districts. Monitoring of jack pine plantations showed that populations of the white pine weevil had declined slightly while damage by the eastern pine shoot borer increased. Although it did not affect commercial forest species, an extensive infestation of leaf beetles caused spectacular discoloration of willow across the region.

The most important disease in 1988 was western gall rust, which caused substantial mortality in young jack pine plantations. An unusual abiotic condition was found on spruce in several districts.

Pheromone trapping to detect the presence of gypsy moth was continued in provincial parks, and captures were made in Kirkland Lake District. Other special projects included experimental pheromone trapping for spruce budworm, detailed surveys of jack pine plantations, monitoring plots established for acid rain studies and surveys for the distribution of pinewood nematode.

The impending retirement of Walter Jansons has necessitated staff changes that have not yet been completed. The assistance and cooperation of Ontario Ministry of Natural Resources (OMNR) and woods industry personnel in all districts are gratefully acknowledged.

The same format was followed in categorizing forest pests as in the 1987 Northern Region report.

### **Major Insects/Diseases**

capable of causing serious injury to or death of living trees or shrubs

### **Minor Insects/Diseases**

capable of causing sporadic or localized injury but not usually a serious threat to living trees or shrubs


### Other Forest Insects/Diseases

These tables provide information on two types of pest:

- (1) those that are of minor importance and have not been known to cause serious damage to forest trees;
- (2) those that are capable of causing serious damage but, because of low populations or for other reasons, did not cause serious damage in 1988.

L.S. MacLeod  
V. Jansons  
S. Payne

Frontispiece



Jack pine with leader killed by white pine weevil (*Pissodes strobi* [Peck ])



Control operation of clipping leaders of jack pine killed by white pine weevil (Island Lake Tree Improvement Area, Chapleau District, July 1988)

# TABLE OF CONTENTS

	Page
INSECTS	
Major Insects	
Spruce Budworm, <i>Choristoneura fumiferana</i> . . . . .	1
(All districts)	
Jack Pine Budworm, <i>Choristoneura pinus pinus</i> . . . . .	1
(Chapleau, Cochrane, Hearst, Kapuskasing, Kirkland Lake and Timmins districts)	
Eastern Pine Shoot Borer, <i>Eucosma gloriola</i> . . . . .	6
(Chapleau, Gogama, Kirkland Lake and Timmins districts)	
Birch Leafminer, <i>Fenusa pusilla</i> . . . . .	7
(All districts)	
Forest Tent Caterpillar, <i>Malacosoma disstria</i> . . . . .	7
(Chapleau, Hearst, Kapuskasing and Kirkland Lake districts)	
Whitespotted Sawyer, <i>Monochamus scutellatus</i> . . . . .	10
(Chapleau District)	
Pine Sawflies, <i>Neodiprion maurus</i> , <i>N. nanulus nanulus</i> and <i>N. virginiana</i> complex . . . . .	10
(Chapleau, Gogama, Hearst and Kirkland Lake districts)	
Swaine Jack Pine Sawfly, <i>Neodiprion swaini</i> . . . . .	11
(Chapleau, Cochrane and Kirkland Lake districts)	
Yellowheaded Spruce Sawfly, <i>Pikonema alaskensis</i> . . . . .	11
(All districts)	
White Pine Weevil, <i>Pissodes strobi</i> . . . . .	12
(Chapleau, Gogama, Kirkland Lake and Timmins districts)	
Larch Sawfly, <i>Pristiphora erichsonii</i> . . . . .	12
(All districts)	
Mountain-ash Sawfly, <i>Pristiphora geniculata</i> . . . . .	14
(Chapleau and Gogama districts)	

(cont'd)

## TABLE OF CONTENTS (cont'd)

	Page
<b>Major Insects (concl.)</b>	
Early Aspen Leafcurler, <i>Pseudexentera oregonana</i> . . . . .	14
(Hearst District)	
Gray Willow Leaf Beetle, <i>Pyrrhalta decora decora</i> . . . . .	14
(All districts)	
<b>Minor Insects</b>	
Jack Pine Tip Beetle, <i>Conophthorus banksianae</i> . . . . .	14
(Chapleau and Gogama districts)	
Other forest insects . . . . .	16
 <b>TREE DISEASES</b>	
<b>Major Diseases</b>	
Armillaria Root Rot, <i>Armillaria mellea</i> . . . . .	21
(All districts)	
Scleroderris Canker, <i>Ascocalyx abietina</i> . . . . .	21
(Chapleau, Gogama, Kirkland Lake and Timmins districts)	
Pine Needle Rust, <i>Coleosporium asterum</i> . . . . .	22
(Chapleau, Gogama, Hearst and Kirkland Lake districts)	
White Pine Blister Rust, <i>Cronartium ribicola</i> . . . . .	22
(Chapleau district)	
Tar Spot Needle Cast, <i>Davisomycella ampla</i> . . . . .	22
(Chapleau, Gogama, Kirkland Lake and Timmins districts)	
Western Gall Rust, <i>Endocronartium harknessii</i> . . . . .	24
(Chapleau, Gogama, Kirkland Lake and Timmins districts)	
Other forest diseases . . . . .	25

(cont'd)

## TABLE OF CONTENTS (concl.)

	Page
ABIOTIC DAMAGE	
Atrophic Growth on Spruce . . . . . (Chapleau, Cochrane, Hearst and Kapuskasing districts)	28
SPECIAL SURVEYS	
Jack Pine Plantations . . . . . (All districts)	30
Surveys of Nurseries . . . . . (Chapleau, Gogama and Kirkland Lake districts)	33
Gypsy Moth Pheromone Traps . . . . . (All districts)	33
Acid Rain National Early Warning System . . . . . (Chapleau, Kapuskasing and Kirkland Lake districts)	34
Jack Pine Seed and Cone Pests . . . . . (Chapleau and Hearst districts)	34
Pinewood Nematode . . . . . (All districts)	35
Climatic Data . . . . .	35



## INSECTS

### Major Insects

#### Spruce Budworm, *Choristoneura fumiferana* (Clem.)

Spruce budworm populations continued to decline in Ontario, with a reduction in the total area of moderate-to-severe defoliation from 7,189,763 ha in 1987 to 5,224,734 ha in 1988. All defoliation was confined to the North Central and Northwestern regions (Fig. 1).

For the second consecutive year ground and aerial detection surveys failed to reveal any damage by the spruce budworm in the Northern Region. Low numbers of larvae were found during third-instar sampling in Hearst District, where only four and seven larvae, respectively, were recovered from sampling stations in Larkin and Frost townships. Similar sampling in Chapleau and Kirkland Lake districts yielded negative results in 1988.

Pheromone trapping was carried out on impact plots and in selected locations, but only small numbers of adults were recovered (Table 1).

A modest increase in adult moth captures was recorded in light trap operations at Remi Lake in Kapuskasing District. A similar situation was found at Chapleau, except for the nights of 8, 9 and 10 July, when 1,256 moths were trapped (Table 2), a fact that suggests a large influx of adults from an unknown source.

Egg-mass sampling was carried out at 35 locations across the region, and results indicate a low potential for extensive damage in 1989 (Table 3).

#### Jack Pine Budworm, *Choristoneura pinus pinus* Free.

Although this insect virtually disappeared from the region in 1987, detection surveys were carried out in jack pine (*Pinus banksiana* Lamb.) plantations and natural stands, particularly in areas that had been infested recently. The only budworm found were two adult moths captured in a light trap operated at Chapleau. Egg-mass surveys were completed at 12 locations in the Chapleau, Cochrane, Hearst, Kapuskasing, Kirkland Lake and Timmins districts; as not a single egg mass was recovered, however, this budworm is not expected to be a problem in 1989.

Table 1. Summary of captures of male spruce budworm moths in pheromone traps in six districts of the region, 1986-1988.

Location (Twp)	Total moths captured		
	1986	1987	1988
<u>Chapleau District</u>			
Peters	99	0	0
Neelands	14	0	0
<u>Cochrane District</u>			
St. John	7	0	2
<u>Gogama District</u>			
Dublin	218	0	1
<u>Hearst District</u>			
Larkin	63	6	1
Frost	55	29	7
<u>Kapuskasing District</u>			
Fauquier	8	1	1
Guilfoyle	41	0	4
<u>Kirkland Lake District</u>			
Maisonville	0	0	0
Pacaud	12	0	0

Table 2. Summary of captures of spruce budworm moths in light traps at two locations in the region, 1987-1988.

Location	Total moths captured	
	1987	1988
<u>Kapuskasing District</u>		
Remi Lake	4	87
<u>Chapleau District</u>		
Chapleau	4	1375

# NORTHWESTERN ONTARIO




Figure 1. Spruce budworm (*Choristoneura fumiferana* [Clem.]).

Areas within which moderate-to-severe  
defoliation occurred in 1988



or •

Table 3. Spruce budworm: Summary of defoliation estimates and egg-mass counts in 1988, and infestation forecasts for 1989.

Location	Host	Estimated defoliation in 1988 (%)	No. of egg masses per 9.29 m <sup>2</sup> of foliage	Infestation forecasts for 1989 <sup>a</sup>	Accumulated damage <sup>b</sup>
<u>Chapleau District</u> (4 locations)					
Neelands Twp - Impact Plot	bF	0	0	0	0
	wS	0	0	0	0
Peters Twp					
- Shoals Prov. Pk	bF	0	42	L-M	+
Reaney Twp					
- Five Mile Prov. Pk	bF	0	13	L	0
<u>Cochrane District</u> (5 locations)					
Clute Twp - OMNR SPA <sup>c</sup>	wS	0	0	0	0
Fournier Twp - OMNR SPA <sup>c</sup>	wS	0	0	0	0
Laughton Twp	bF	0	0	0	9
Marathon Twp	bF	0	0	0	9
St. John Twp - Stand 177	bF	0	0	0	9
<u>Gogama District</u> (2 locations)					
Dublin Twp	bF	0	0	0	0
Garvey Twp	bF	0	26	L-M	0
<u>Hearst District</u> (16 locations)					
Elgie Twp	bF	0	0	0	9
Franz Twp	bF	0	0	0	9
Frost Twp					
- Nagagamisis Prov. Pk Campground	bF	0	0	0	9
- Impact Plot	bF	0	0	0	9
	wS	0	35	L-M	9
	bS	0	0	0	9
Pushimi Twp					
- Fushimi Prov. Pk	bF	0	30	L-M	9
Kohler Twp	bF	0	0	0	9

(cont'd)

Table 3. Spruce budworm: Summary of defoliation estimates and egg-mass counts in 1988, and infestation forecasts for 1989 (concl.).

Location	Host	Estimated defoliation in 1988 (%)	No. of egg masses per 9.29 m <sup>2</sup> of foliage	Infestation forecasts for 1989 <sup>a</sup>	Accumulated damage <sup>b</sup>
<u>Hearst District (cont'd)</u> (16 locations)					
Larkin Twp					
- Impact Plot	bF	0	0	0	9
	vS	0	0	0	9
	bS	0	0	0	9
Lessard Twp - Stand 341	bF	0	34	L-M	9
McEwing Twp	bF	0	0	0	9
Rogers Twp					
- Plantation 26A	bF	0	0	0	9
	vS	0	0	0	9
Studholme Twp					
- (Plantation 37)	vS	0	0	0	0
<u>Kapuskasing District</u> (3 locations)					
Cumming Twp	vs	0	0	0	9
Fauquier Twp					
- René Brunelle Prov. Pk	bF	0	0	0	9
Guilfoyle Twp	bF	0	0	0	9
<u>Kirkland Lake District</u> (3 locations)					
Maisonville Twp	bF	0	0	0	+
	bS	0	0	0	+
Pacaud Twp	bF	0	0	0	+
<u>Timmins District</u> (2 locations)					
Hassard Twp	bF	0	0	0	+
Thomas Twp	vS	0	0	0	3

<sup>a</sup> S = severe, M = moderate, L = light, 0 = nil

<sup>b</sup> 0 = undamaged, 3 = severe damage (60 to 80% total defoliation, 3 to 5 seasons of severe defoliation, will recover), 9 = less than 25% of stand dead (no significant defoliation, 0-25%, for several years), + = 25 to 50% of stand dead (no significant defoliation for several years)

<sup>c</sup> SPA = Seed Production Area

Eastern Pine Shoot Borer, *Eucosma gloriola* Heinr.

Leader mortality caused by the eastern pine shoot borer was found in most jack pine plantations examined in the southern districts of the region (see photo page). In the Chapleau District, there was 15.3% leader damage at the Island Lake Tree Improvement Area, and in Copperfield Township, 10% of the leaders were killed. Quantitative sampling also indicated a general increase in population levels at several locations in the Kirkland Lake and Timmins districts (Table 4).

Table 4. Summary of leader damage by the eastern pine shoot borer in four districts of the region in 1987 and 1988 (counts based on the examination of 150 randomly selected jack pine trees at each location).

Location (Twp)	Avg ht of trees (m)	Estimated no. of trees per ha	Estimated area affected (ha)	Leaders killed (%)	
				1987	1988
<u>Chapleau District</u>					
Marshall	2.4	1500	25	0.0	0.7
Island Lake TIA <sup>a</sup>	1.3	2500	20	3.3	15.3
Copperfield	2.2	1500	68	0.0	10.0
Dalmas	2.3	1500	25	4.0	3.3
Hutcheon	1.8	1600	10	0.0	0.7
McNaught	1.4	1500	50	-	2.0
<u>Gogama District</u>					
Sheard	2.2	2500	200	0.0	2.0
<u>Timmins District</u>					
Robb	3.0	2500	100	2.0	3.3
Macklem	1.8	3500	50	3.0	8.6
Kenogaming	3.5	3500	50		0.3
Carscallen	2.5	4000	50	-	12.0
<u>Kirkland Lake District</u>					
Holmes	2.8	4000	50	-	9.3
Charters	1.0	4000	50	-	5.3
Chown	0.8	4000	50	-	4.0
Gross	2.1	4000	50	1.3	3.3

<sup>a</sup> TIA = Tree Improvement Area

**Birch Leafminer, *Fenusa pusilla* (Lep.)**

High populations of the birch leafminer were again found in most districts of the region. Foliage discoloration and defoliation by this introduced pest were particularly conspicuous on ornamental birches (*Betula* spp.) through urban areas, on open-grown white birch (*Betula papyrifera* Marsh.) in rural situations, and in lightly stocked forest stands.

In the towns of Chapleau and Gogama every birch tree examined was infested and 99% of the foliage was mined. Severe mining, frequently in 90% of the foliage, was common in the towns of Iroquois Falls, Cochrane, Kapuskasing and Hearst. In forested areas moderate-to-severe damage was observed on fringe trees along Highway 631 in Arnott, McEwing, Frost and Elgie townships, Hearst District, along several access roads in Kapuskasing District, and in the rural area north of Cochrane in Cochrane District.

In Kirkland Lake District, 50% foliar damage was common in pockets ranging from 0.5 to 2 ha in the Elk Lake, Earlton, Charlton, Englehart, Kirkland Lake, Larder Lake and Matachewan areas. Moderate-to-severe defoliation of ornamental trees was recorded in South Porcupine and Timmins in Timmins District.

**Forest Tent Caterpillar, *Malacosoma disstria* Hbn.**

Several changes occurred in the forest tent caterpillar situation in the region in 1988. In Kirkland Lake District the outbreak subsided abruptly, and although a few colonies of caterpillars were found through poplar (*Populus* spp.) stands in the spring and early summer, no defoliation was observed during ground and aerial surveys.

In the northern part of Hearst District, a new infestation totaling 9,930 ha caused moderate-to-severe defoliation on narrow bands of trembling aspen (*Populus tremuloides* Michx.) growing along the Nagagami, Kenogami, Kabinakagami and Pitukani rivers near their junction at the abandoned village of Mammamattawa. As well, three small pockets totaling 620 ha were mapped in Studholme, Stoddart and Kendall townships (Fig. 2). The insect was not found in other districts of the region.

Monitoring of light traps operated at Chapleau and at Remi Lake disclosed caterpillar moth captures of 395 and 58, respectively. This represents an increase over 1987, when 72 and 12 moths were taken, respectively.

Egg-band counts in the autumn of 1988 indicate that the infestation near Hearst will increase substantially in extent and intensity in 1989 (Table 5). Inaccessibility and weather problems prevented egg surveys in the Mammamattawa segment of the infestation but that infestation is also expected to expand considerably in 1989 (Fig. 3).

## NORTHERN REGION




Figure 2. Forest tent caterpillar (*Malacosoma disstria* Hbn.).

Areas within which moderate-to-severe defoliation  
occurred in 1988



# NORTHERN REGION




Figure 3. Forest tent caterpillar (*Malacosoma disstria* Hbn.).

Locations of egg-band samples taken in 1988 and infestation forecasts for 1989:

Negative	○
Light	◐
Moderate to heavy	●

Table 5. Summary of forest tent caterpillar egg-band counts on trembling aspen in two districts of the region in 1988 and infestation forecasts for 1989.

Location (Twp)	Avg DBH of trees (cm)	No. of trees sampled	Total no. of egg bands	Infestation forecasts for 1989
<u>Hearst District</u>				
Kendall	8	2	16	heavy
Studholme	11	3	32	heavy
Rogers	16	3	0	nil
238	31	3	1	light
<u>Kirkland Lake District</u>				
Tyrell	16	3	0	nil
Milner	20	3	0	nil
Harley	15	3	0	nil
Casey	17	3	0	nil

**Whitespotted Sawyer, *Monochamus scutellatus* (Say)**

Feeding by adult sawyer beetles caused appreciable damage in one jack pine plantation in Delaney Township, Chapleau District (see photo page). Defoliation was concentrated in a 2-ha area adjacent to a jack pine log deck. In the plantation, defoliation of 99% was recorded on 1.7-m trees growing within 15 m of the log piles. Beyond 15 m the damage became progressively lighter and the total area infested did not exceed 8 ha.

**Pine Sawflies, *Neodiprion maurus* Roh., *N. nanulus nanulus* Schedl., and *N. virginiana* complex**

Although these sawflies did not cause serious defoliation in the region they were found more frequently than in previous years. Colonies of *Neodiprion virginiana* were common in jack pine stands throughout Chapleau, Gogama and Kirkland Lake districts. Single trees and small groups of trees along the fringes of stands sustained up to 30% defoliation at many locations, particularly in Margaret Township, Chapleau District and in Skead, Catherine, Mickle, Kimberly and Otto townships in Kirkland Lake District. Scattered colonies of *N. nanulus nanulus* were found on jack pine trees in Frost Township, Hearst District and at several points in the southern part of Kirkland Lake District. A single colony of the relatively rare sawfly *N. maurus* was found at Watabeag Lake in Nordica Township, Kirkland Lake District.

**Swaine Jack Pine Sawfly, *Neodiprion swainei* Midd.**

The Swaine jack pine sawfly infestation in the Elk Lake Management Unit in Kirkland Lake and Temagami districts peaked in 1981 when an area of 5,699 ha of jack pine was moderately to severely defoliated. From 1982 to 1984 a steady downward trend in population levels was recorded and the infestation subsided completely by 1985.

On 22 August 1988 a helicopter survey was made over the former infestation in the Lady Evelyn-Banks-Makobe-Alexander lakes area and light defoliation was mapped in two pockets of jack pine totaling 200 ha between Trethewey and Wolfskin lakes in the northwestern part of Trethewey Township. Subsequent ground checks revealed scattered colonies of the sawfly in jack pine plantations in Brewster, Gamble and McGiffin townships, Temagami District; in Corkill, Cane and Pacaud townships, Kirkland Lake District; in Caouette Township, Chapleau District; and in Greenwater Provincial Park, Cochrane District.

Moderate-to-severe defoliation (approximately 50%) recurred on Island 127 on the northeast arm of Lake Temagami, where a small pocket (0.2 ha) of infestation has persisted for several years. Scattered colonies of the sawfly were also found on islands and in shoreline stands at many other points on Lake Temagami.

**Yellowheaded Spruce Sawfly, *Pikonema alaskensis* (Roh.)**

High numbers of the yellowheaded spruce sawfly and resultant damage were reported from several districts of the region. Windbreaks, snow hedges, plantations and white spruce (*Picea glauca* [Moench] Voss) and black spruce (*Picea mariana* [Mill.] B.S.P.) trees in open-growing situations along roadsides and in old fields and cutovers sustained moderate-to-severe defoliation at many locations.

In the town of Chapleau approximately 40% of the ornamental trees experienced 10-75% defoliation. In Wakami Provincial Park defoliation was approximately 40% on 100% of the 1- to 2-m spruce trees throughout the park. Ornamental white spruce trees at the entrance to Ivanhoe Provincial Park were also defoliated, with damage ranging from 10 to 35%. In Battersby Township, Gogama District some damage occurred in a mixed pine and spruce plantation but current defoliation did not exceed 10%.

In Kirkland Lake District the sawfly was found infesting one-year-old black spruce trees in the OMNR seed orchard in Chamberlain Township. Windbreaks at the Swastika Tree Nursery were sprayed to control defoliation, and spraying was also carried out by the Ontario Ministry of Transportation and Communications on snow hedges along Highway 11 between New Liskeard and Englehart. The most severe damage in the district was recorded in the Ramore-Shillington-Matheson-Monteith area, where defoliation ranged from 10 to 80%.

Moderate-to-severe defoliation on trees averaging 2 m in open-growing situations occurred at numerous points in Timmins, Cochrane and Kapuskasing districts. Defoliation ranged from 10 to 90% on black and white spruce in Mowbray and Howells townships, Kapuskasing District and on 2.5-m white spruce trees at several points in Fournier Township, Cochrane District. Incidence and damage levels decreased on 1- to 2-m white spruce trees in René Brunelle Provincial Park at Remi Lake in Kapuskasing District, and little damage resulted.

#### White Pine Weevil, *Pissodes strobi* (Peck)

Relatively high leader mortality in some jack pine plantations in several districts of the region, together with concern expressed by woods personnel of the E.B. Eddy Company, prompted a detailed survey to assess damage by this pest in 1988. In April, FIDS crews working from E.B. Eddy's Camp 303 conducted surveys of jack pine plantations in Invergarry and Vrooman townships, Gogama District. Forty semi-permanent plots were established and damage incurred by the weevil and other agents in 1987 was recorded. These plots were retallied in the autumn of 1988 to provide a comparison of damage by the various pests (Table 6).

Results of quantitative sampling for the white pine weevil in other plantations of the region are shown in Table 7 (see also Frontispiece).

#### Larch Sawfly, *Pristiphora erichsonii* (Htg.)

In 1987 intensive surveys of tamarack (*Larix laricina* [Du Roi] K. Koch) stands failed to reveal any evidence of this sawfly in the region. Stands were monitored closely in 1988 and several colonies were recovered from Fauquier and Macvicar townships, Kapuskasing District. Curled shoots with egg slits and a few branches showing light defoliation were observed at one location in Thorneloe Township, Timmins District but no sawflies were found here or in other districts of the region.

Table 6. Summary of leader damage in jack pine plantations in two townships of Gogama District, 1987-1988 (counts based on the examination of 150 randomly selected trees at each location).

Location (Twp)	Year	White pine weevil	Eastern pine shoot borer	Jack pine tip beetle	Unknown
Invergarry	1987	8.0	3.6	1.4	4.5
	1988	3.6	5.5	0.8	9.4
Vrooman	1987	4.4	0.9	0.2	2.8
	1988	1.1	0.9	0.4	3.3

Table 7. Summary of damage caused by the white pine weevil in four districts of the region in 1987 and 1988 (counts based on the examination of 150 randomly selected jack pine trees at each location).

Location (Twp)	Avg ht of trees (m)	Estimated no. of trees per ha	Estimated area affected (ha)	Leaders killed (%)	
				1987	1988
<u>Chapleau District</u>					
Marshall	2.4	1500	25	10.7	6.0
Island Lake TIA	1.3	2500	20	10.7	8.0
Copperfield	2.2	1500	68	0.7	2.7
Hutcheon	1.8	1600	10	14.7	4.7
De Gaulle	1.8	2500	10	-	2.0
Langlois	2.1	2500	15	20.0	3.3
Caverly	2.1	2500	50	-	8.0
Dalmas	2.3	1500	25	19.3	16.0
McNaught	1.4	1500	50	-	11.3
Esther	2.4	1500	10	-	2.7
Edith	3.1	1500	25	-	4.7
<u>Gogama District</u>					
Sheard	2.2	2500	200	22.3	15.7
MacMurchy	1.5	1500	20	-	2.7
Benneweis	2.7	1500	10	-	4.7
<u>Timmins District</u>					
Macklem	1.8	3500	50	12.6	9.3
Kenogaming	3.5	3500	50	-	5.3
Carscallen	2.5	4000	50	-	9.3
<u>Kirkland Lake District</u>					
Holmes	2.8	4000	50	-	20.6
Charters	1.0	4000	50	-	31.3
Chown	0.8	4000	50	-	21.3
Gross	2.1	4000	50	20.6	14.0
Arnold	0.6	4000	50	-	6.6

**Mountain-ash Sawfly, *Pristiphora geniculata* (Htg.)**

Populations of this sawfly remained relatively low throughout the region. In the town of Chapleau defoliation of ornamental trees did not exceed 30%. Heavier damage was recorded at Ivanhoe Provincial Park, where defoliation of mountain-ash (*Sorbus americana* Marsh.) trees averaged 60% (see photo page). In Battersby Township, Gogama District, 20% of the trees examined sustained foliar damage of 30% but in all other districts the sawfly was rarely found and little damage to either urban or forest trees was observed.

**Early Aspen Leafcurler, *Pseudexentera oregonana* (Wlsm.)**

High numbers of this leafroller recurred at several locations in the Hearst District in the northwestern part of the region (Fig. 4). Moderate-to-severe defoliation of mature trembling aspen stands over a total of 7,500 ha was mapped along the Kabinakagami, Nagagami and Kenogami rivers in the northern part of the district and along the Ridge River east of Indian Reserve #66 (see photo page). Smaller areas of similar damage occurred on scattered aspen stands in Township 238 and Rogers Township. No damage by the insect was reported from other districts of the region.

**Grey Willow Leaf Beetle, *Pyrrhalta decora decora* (Say)**

This insect has been reported in epidemic numbers on willow (*Salix* spp.) and poplar in Newfoundland, western Ontario and the prairie provinces. The tiny adult beetles emerge from hibernation in late May or early June and feed on willow and poplar. After mating, the females lay eggs at the bases of willow bushes. After hatching the larvae move up to feed on the underside of leaves, which they then skeletonize. Larvae complete their feeding about mid-August in Ontario, drop to the soil and pupate. The adults emerge in about seven days and feed on willow and poplar before seeking hibernation sites in protected places.

In 1988, severe skeletonizing of hundreds of hectares of willow throughout Chapleau, Gogama, Kirkland Lake and Timmins districts precipitated numerous inquiries from concerned property owners. In late August light-to-moderate defoliation of trembling aspen was caused by adult feeding at many locations in Kirkland Lake District.

**Minor Insects**

**Jack Pine Tip Beetle, *Conophthorus banksianae* McP.**

These beetles bore into the tips of jack pine twigs in May and leave a characteristic pitch-encrusted entrance hole. In early June egg

## NORTHERN REGION




Figure 4. Early aspen leafcurler (*Pseudexentera oregonana* [Wlsm.]).

Areas within which defoliation of trembling  
aspen occurred in 1988



tunnels constructed by the females cause the new growth to wilt and die (see photo page). Widely distributed through jack pine plantations in the region, the principal damage occurs when the leaders of young trees are destroyed, causing malformation of the main stem (Table 8).

Table 8. Summary of leader damage by the jack pine tip beetle in two districts of the region in 1988 (based on the examination of 150 randomly selected jack pine trees at each location).

Location (Twp)	Avg ht of trees (m)	Estimated area affected (m)	Trees attacked (%)	Leaders killed (%)
<u>Chapleau District</u>				
Hutcheon	1.6	25	50.0	3.0
Langlois	1.3	15	35.3	6.7
<u>Gogama District</u>				
Battersby	2.0	100	60.0	2.0
Invergarry	2.0	50	8.0	1.0

Table 9. Other forest insects.

Insect	Host(s)	Remarks
<i>Aceria</i> sp. nr. <i>dispar</i> (Nal.) Aspen leaf mite	tA	common throughout Timmins and Kirkland Lake districts; particularly conspicuous in stands with foliar damage by other insects
<i>Acrobasis betulella</i> Hlst. Birch tubemaker	wB	less than 5% defoliation in the majority of stands examined in Chapleau, Timmins and Kirkland Lake districts
<i>Actebia fennica</i> (Tausch.) Black army cutworm	-	21 and 31 adults captured on prescribed burn sites in Machin Twp, Kapuskasing District and in Arnott Twp, Hearst District, respectively

(cont'd)



Table 9. Other forest insects (cont'd).

Insect	Host(s)	Remarks
<i>Agromyza</i> sp. nr. <i>alni betulae</i>	wB	about 2% leaf mining on semimature trees in Reeves Twp SPA, Chapleau District
<i>Altica ambiens alni</i> Harr. Alder flea beetle	Al	75% defoliation of alder ( <i>Alnus</i> spp.) shrubs in Hill and Caouette twps, Chapleau District, in Greenwater Provincial Park, Cochrane District and in Munro, McCool and Michaud twps, Kirkland Lake District
<i>Anacampsis niveopulvella</i> (Chamb.) Paleheaded aspen leafroller	tA	less than 10% defoliation of single trees scattered throughout Chapleau and Kirkland Lake districts
<i>Aphrophora cribrata</i> (Wlk.) Pine spittlebug and <i>Aphrophora saratogensis</i> (Fitch) Saratoga spittlebug	jP	large numbers of spittle masses at many points in the southern part of the region; small numbers of nymphs found at Iroquois Falls, Cochrane District
<i>Archips cerasivorana</i> (Fitch) Uglynest caterpillar	Ch	numerous nests in the vicinity of Charlton, Englehart and New Liskeard, Kirkland Lake District; small numbers in Strathearn Twp, Chapleau District
<i>Arge clavicornis</i> (F.) Willow argid	Al	10% defoliation of roadside alder in Strathearn Twp, Chapleau District
<i>Argyresthia laricella</i> Kft. Larch shoot moth	tL	5% of branch tips killed on scattered trees in Calder Twp, Cochrane District
<i>Campaea perlata</i> (Gn.) Fringed looper	tA	5% defoliation of a semi-mature stand, Green Twp, Chapleau District

(cont'd)

Table 9. Other forest insects (cont'd).

Insect	Host(s)	Remarks
<i>Cecidomyia resinicola</i> (O.S) Jack pine resin midge	jP	10-15% shoot mortality in scattered plantations in Chapleau and Kirkland Lake districts
<i>Eupareophora parca</i> (Cress.) Spiny ash sawfly	bAs	90% of stands in Chapleau and Gogama districts affected with about 30% defoliation; less than 5% defoliation in a few stands in Kirkland Lake District
<i>Fenusa dohrnii</i> (Tisch.) European alder leafminer	Al	about 10% foliage mined at several points in Holmes and Nordica twps, Kirkland Lake District
<i>Gonioctena americana</i> (Schaeff.) American aspen beetle	tA	90% defoliation of 2- to 3-m trees growing in jack pine plantations at several points in Chapleau District; low levels throughout Kirkland Lake District
<i>Hyphantria cunea</i> (Drury) Fall webworm	deciduous	5-10% defoliation of willow and alder trees at many points in the vicinity of Englehart, Tomstown, Hilliardton and Belle Vallee, Kirkland Lake District
<i>Malacosoma californicum pluviale</i> (Dyar) Northern tent caterpillar	deciduous	Populations increased markedly in Chapleau and Gogama districts. Tents were common along bush roads and in cut-over stands in Kirkland Lake District.
<i>Petrova albicapitana</i> (Bsk.) Northern pitch twig moth	jP	found in most stands examined, but incidence usually less than 3%, and little damage observed


(cont'd)

Table 9. Other forest insects (concl.).

Insect	Host(s)	Remarks
<i>Phyllonorycter nipigon</i> (Free.) Balsam poplar leafblotch miner	bPo	small pockets of severe mining (75%) at many points in Timmins and Kirkland Lake districts
<i>Phyllonorycter ontario</i> (Free.) Aspen leafblotch miner	tA	mining of up to 80% of foliage at many points in the region, but infested areas small
<i>Pleroneura brunneicornis</i> Roh. Balsam shootboring sawfly	bF	widespread but in low numbers throughout Kirkland Lake and Timmins districts
<i>Profenusa thomsonii</i> (Konow) Ambermarked birch leafminer	wB	occasional mined leaves in Kirkland Lake and Timmins districts; usually less than 5% of the foliage affected
<i>Proteoteras moffatiana</i> Fern. Maple shoot borer	sM	less than 5% foliar damage to mature trees in a 2-ha stand in Deans Twp, Chapleau District
<i>Rhabdophaga swainei</i> Felt Spruce bud midge	bS	10-15% bud damage on open-growing trees, Dempsey Twp, Cochrane District
<i>Tetralopha aplastella</i> (Hlst.) Aspen webworm	tA	approximately 5% defoliation in most stands examined in Timmins and Kirkland Lake districts
<i>Toumeyella parvicornis</i> (Ckll.) Pine tortoise scale	jP	single trees and small groups of trees heavily infested at many points throughout the southern part of the region
<i>Trichiocampus gregarius</i> (Dyar) Hairy poplar sawfly	tA	approximately 10% defoliation of semimature trees at one point in Lloyd Twp, Chapleau District




Jack pine with bark  
stripped by whitespotted  
sawyer beetle (*Monochamus*  
*scutellatus* [Say]) in  
Delaney Twp, Chapleau  
District, June 1988




Adult sawyer beetles feeding on jack pine



Abiotic condition causing  
atrophic growth of white  
spruce shoots in 1988



Club top on jack pine caused  
by stem gall of western gall  
rust (*Endocronartium*  
*harknessii* [J.P. Moore] Y.  
Hirats.) (The gall has not  
yet begun to fruit.)



Jack pine leader killed  
by jack pine tip beetle  
(*Conophthorus banksianae*  
McP.) (Note exit hole  
at base of shoot.)


Young jack pine with leader  
broken by eastern pine shoot  
borer (*Eucosma gloriola* Heinr.)







Aerial photo of severe defoliation by early aspen leafroller (*Pseudexentera oregonana* [Wlsm.]) along Ridge River, Hearst District



Ornamental mountain ash defoliated by mountain-ash sawfly (*Pristiphora geniculata* [Htg.]) in Ivanhoe Provincial Park, Chapleau District, July 1988

## TREE DISEASES

### Major Diseases

Armillaria Root Rot, *Armillaria mellea* (Vahl:Fr.) Kummer

As in previous years, trace levels of this root rot were found throughout the region. The majority of plantations surveyed contained this disease but mortality was not extensive. Damage levels in the sample plots listed in Table 10 are indicative of the levels found throughout the region.

Table 10. Summary of mortality caused by *Armillaria* root rot on jack pine in four districts of the region in 1988 (counts based on the examination of 150 randomly selected trees at each location).

Location (Twp)	Avg ht of trees (m)	Estimated no. of trees/ha	Area affected (ha)	Current mortality (%)
<u>Cochrane District</u>				
Freele Twp	1.8	1,600	3	0.7
<u>Gogama District</u>				
Invergarry	1.0	1,600	10	0.7
<u>Kirkland Lake District</u>				
Morrisette	1.2	3,800	15	1.9
Pense	0.9	4,000	6	0.7
Catherine	1.3	3,500	3	2.0
Corkill	0.8	3,500	5	1.9
Arnold	0.8	3,500	10	0.7
<u>Timmins District</u>				
Macklem	0.7	3,300	3	2.6

Scleroderris Canker, *Ascocalyx abietina* (Lagerb.) Schläpfer-Bernhard

Surveys were conducted in red pine (*Pinus resinosa* Ait.) and jack pine plantations throughout the region to locate the presence of *Scleroderris* canker. Surveys revealed trace levels of the disease in Chapleau, Gogama, Kirkland Lake and Timmins districts. These results were consistent with the previous year's results, an indication of either no increase or a decrease in the incidence of this disease.



There are two races of this disease, the native and the European. The European race is much more serious, as it can kill trees of all sizes, and in one season. The native race kills only trees that are less than 2 m high. All samples of *Scleroderris* canker found were submitted to the Disease Survey staff at the Great Lakes Forestry Centre for race identification. All tests for the European race were negative in 1988.

**Pine Needle Rust, *Coleosporium asterum* (Dietel) Sydow**

In 1988, the incidence of this foliage disease of jack pine increased throughout the entire region. Light damage was common in the majority of plantations throughout the southern portion of the region (Table 11). Severe damage was found in several areas, the heaviest occurring in Reaney Township, Chapleau District, where 100% of the trees were affected and there was an average of 66% foliar damage. This disease causes the premature loss of the previous year's foliage. Light damage has little impact on the tree. However, severe infection can kill small trees and cause a reduction in growth of trees of sapling size.

**White Pine Blister Rust, *Cronartium ribicola* J.C. Fisher**

This disease is common on mature white pine (*Pinus strobus* L.) in the southern part of the region. Trees with tops killed by blister rust are a common sight when one is traveling through some districts. It is difficult to assess damage to mature trees because they are so widely scattered, but a reasonable estimate would be that 15% of the trees are affected.

In Chapleau District, recently planted trees were affected in four townships (Table 12). In all cases, the trees were severely infected, with the infection having reached the stem. This infection will eventually kill the trees. In the areas evaluated, it was not possible to do a normal 150-tree survey; therefore, a 100% evaluation was completed.

**Tar Spot Needle Cast, *Davisonmycella ampla* (J. Davis) Darker**

Occurrence of this early-season foliage disease of jack pine was generally low throughout the region. Needle cast causes the previous year's foliage to discolor and to fall off prematurely. In years of heavy defoliation, trees are significantly weakened and show a reduction in growth. Although damage was generally light in most of the region, a few pockets of moderate infection were surveyed (Table 13).

Table 11. Summary of damage caused by needle rust of jack pine in four districts of the region in 1988 (counts based on the examination of 150 randomly selected trees at each location).

Location (Twp)	Avg ht of trees (m)	Estimated no. of trees per ha	Estimated area affected (ha)	Trees affected (%)	Foliar damage (%)
<u>Chapleau District</u>					
Copperfield	0.4	4,444	100	75	5
Reaney	0.3	4,444	25	100	66
Marshall	1.5	2,500	30	99	12
D'Arcy	1.3	2,500	60	100	22
Chappise	2.1	1,600	10	60	10
Nimitz	1.0	2,500	134	13	6
Smuts	0.4	4,444	30	100	15
Warren	1.0	2,500	10	100	33
<u>Gogama District</u>					
Garibaldi	1.2	2,500	50	100	15
Invergarry	0.5	2,500	60	75	10
MacMurchy	1.5	2,500	25	100	22
<u>Hearst District</u>					
Cross	3.5	2,200	220	100	5
Larkin	0.8	4,500	5	100	5
<u>Kirkland Lake District</u>					
Tyrell	1.7	2,800	5	90	15
Ingram	7.0	2,500	10	3	2
Arnold	0.8	4,444	10	3	1

Table 12. Summary of damage caused by white pine blister rust in Chapleau District in 1988.

Location (Twp)	Avg ht (m)	No. of trees examined	No. of trees affected	Trees affected (%)
Gilliland	0.8	60	3	5
Kosny	0.7	25	4	16
Green	1.0	25	1	4
Neill	0.8	22	6	27

Table 13. Summary of damage caused by needle cast of jack pine in four districts of the region in 1988 (counts based on the examination of 150 randomly selected trees at each location).

Location (Twp)	Avg ht of trees (m)	Estimated no. of trees per ha	Estimated area affected (ha)	Trees affected (%)	Defolia- tion level (%)
<u>Chapleau District</u>					
Nimitz	1.0	2,500	134	40.7	12
DeGaulle	1.5	2,500	10	51.3	10
Hutcheon	1.6	2,500	20	21.3	6
Chappise	2.1	2,500	23	23.3	13
Neelands	1.2	2,500	5	42.0	7
<u>Gogama District</u>					
Invergarry	1.0	2,500	5	24.7	11
Garibaldi	1.5	2,500	50	40.0	5
<u>Kirkland Lake District</u>					
Tyrell	3.0	4,500	100	1.0	10
Cane	8.0	3,500	50	7.0	10
Ingram	6.0	3,500	10	6.0	10
Gauthier	5.0	2,500	2	2.0	10
Gross	2.0	3,500	50	5.0	10
Beauchamp	6.0	2,000	50	2.0	10
<u>Timmins District</u>					
Robb	3.0	4,500	100	26.0	10

Western Gall Rust, *Endocronartium harknessii* (J.P. Moore) Y. Hirats.

Areas with high infection levels in 1987 were again surveyed in 1988 in Chapleau, Gogama and Kirkland Lake districts (Table 14), and several new areas of severe infection of jack pine were evaluated. As the galls are located on the stem, many of the infected trees will probably die eventually. Stem galls cause abnormal growth of young trees (see photo page), affecting their form.

Infected trees at Island Lake TIA were pulled out in 1987 in a control operation that proved to be very successful. In 1987, 18% of the trees were severely infested with stem galls. After the control operation, fewer than 1% of the trees were infected.

Table 14. Summary of damage by western gall rust in four districts of the region in 1988 (counts based on the examination of 150 jack pine trees at each location).

Location (Twp)	Avg ht of trees (m)	Estimated no of trees per ha	Estimated no. of ha infected	Trees affected (%)	Trees affected severely (%)
<u>Chapleau District</u>					
Neelands	1.2	4,444	5	37.3	27.3
DeGaulle	1.5	2,500	10	39.3	19.3
Caverly	0.4	4,444	5	30.7	29.3
Chappise	2.2	2,500	25	12.7	0
Copperfield-south	0.5	2,500	5	14.0	14.0
Copperfield-north	0.4	4,444	100	35.3	35.3
Hutcheon	1.6	2,500	20	12.7	4.7
Nimitz	1.0	2,500	135	10.7	6.7
Reaney	1.0	2,500	30	2.0	2.0
<u>Gogama District</u>					
MacMurchy	2.0	2,500	40	8.0	8.0
Invergarry	1.0	2,500	15	9.0	7.6
<u>Kirkland Lake District</u>					
Morrisette	1.0	4,500	50	0.6	0
Gauthier	5.0	2,500	2	1.3	0
Gross	2.0	3,500	50	0.6	0
Ben Nevis	1.0	4,500	10	32.6	32.6
<u>Timmins District</u>					
Robb	3.0	4,500	100	3.3	0.6
Thornloe	4.0	2,500	4	1.3	0
Sewell	4.0	2,500	2	3.3	0
German	4.0	1,500	2	0.6	0

Table 15. Other forest diseases.

Organism	Host(s)	Remarks
<i>Blumeriella jaapii</i> (Rehm) v. Arx Shot-hole	Ch	80-90% foliar discoloration common on shrubs near the town of Chapleau, Chapleau District
<i>Chrysomyxa ledi</i> (Alb. & Schwein) de Bary var. <i>ledi</i> Spruce needle rust	bS	1-2% foliar damage in many areas in the region

(cont'd)

Table 15. Other forest diseases (cont'd).

Organism	Host(s)	Remarks
<i>Chrysomyxa pirolata</i> (Körn.) Winter Spruce cone rust	wS	1-2% of the cones damaged on a few scattered trees at Opasatika TIA, Kapuskasing District
<i>Ciborinia whetzellii</i> (Seaver) Seaver Ink spot of aspen	tA	10% foliar damage to young road-side aspen in Lloyd Twp, Chapleau District; 25% foliar damage to 100% semimature trees in a 0.5-ha clearing in Hellyer Twp, Chapleau District
<i>Cladosporium aecidiicola</i> Thüm.	bS	parasite of needle rust found on small scattered plantings in Abbott Twp, Kapuskasing District
<i>Isthmiella faullii</i> (Darker) Darker Needle cast	bF	10-15% loss of old foliage at scattered locations in central Chapleau District
<i>Leucostoma nivea</i> (Hoffm.:Fr.) Höhnelt Cytospora canker	hybrid poplar	a few dead trees at Swastika Tree Nursery, Kirkland Lake District
<i>Linospora tetraspora</i> G.E. Thompson Linospora leaf blight	bPo	pockets of 75-80% foliar damage throughout the northern part of the region
<i>Lophodermella concolor</i> (Dean) Darker Needle cast	jP	90% foliar damage to individual trees in Kipling Twp, Kapuskasing District and in Corkill Twp, Kirkland Lake District
<i>Lophodermium</i> sp. Needle cast	jP	1-2% defoliation common in Island Lake TIA, Chapleau District
<i>Melampsora medusae</i> Thüm. Larch-poplar rust	hybrid poplar	75% of foliage damaged in a compartment of hybrid poplar, Bonner Centre, Kapuskasing District
<i>Melampsora</i> sp. Willow rust	W	100% of leaves infected on bushes in Shoals Prov. Pk, Chapleau District

(cont'd)

Table 15. Other forest diseases (concl.).

Organisms	Host(s)	Remarks
<b>Sphaeropsis sapinea</b> (Fr.) Dyko & B. Sutton Tip blight	jP	several trees with stem cankers in Ben Nevis Twp, Kirkland Lake District
<b>Uredinopsis sp.</b> Needle rust	bF	100% of examined regeneration trees less than 1.5 m tall infected in Wakami Prov. Pk, Chapleau District, with an average of 25% defoliation
<b>Valsa friesii</b> (Duby) Fuckel Dieback	bF	scattered 1-m trees with branch mortality in Dublin Tp, Gogama District
<b>Valsa sordida</b> Nitschke Cytospora canker	tA	one tree killed in Chown Twp, Kirkland Lake District
<b>Venturia macularis</b> (Fr.) E. Müller & v. Arx Shoot blight	tA	widely distributed in aspen regeneration throughout the region, but damage generally insignificant

## ABIOTIC DAMAGE

### Atrophic Growth on Spruce

There was a high incidence of bud failure on young white spruce and black spruce in northern Chapleau District and central Hearst, Kapuskasing and Cochrane districts (Fig. 5). Most severely affected were plantations of open-growing 1- to 3-m trees, on which up to 100% of the buds were affected in the top and mid-crown sections (see photo page). However, below the 60- to 70-cm level, shoot development was normal. It is interesting to note that this damage was random, with one tree affected and a tree 1 m away unaffected. In most plantations surveyed, roughly two thirds of the trees were affected (Table 16). The reason for this damage is undetermined, but the likely cause was unusual weather during the winter of 1987-1988.

Table 16. Summary of abiotic damage to spruce in Chapleau and Hearst districts in 1988.

Location (Twp)	Host <sup>a</sup>	Avg ht (m)	Trees affected (%)
<u>Chapleau District</u>			
Gilliland	wS	1.2	75
Lipsett	wS	1.6	60
Panet	wS	1.0	70
<u>Kapuskasing District</u>			
Bonner Centre	bS	2.3	65

<sup>a</sup> wS = white spruce, bS = black spruce

# NORTHERN REGION




Figure 5. Atrophic growth on white spruce and black spruce in 1988.

Area within which atrophic growth occurred



## Jack Pine Plantations

A special survey was conducted to assess the impact of insect and disease problems in 18 jack pine plantations in the region (Table 17). Stand selection was from three height classes: 0.5-2 m, 2-6 m, and >6 m. Six of the areas examined in the current survey were the 300-tree semipermanent plots established in 1982. Each of the other 12 stands of jack pine consisted of a 150-tree sample. Two visits were made to each plot during the periods 6-24 June and 1-31 August. The stands were checked for the presence of the following insects and diseases (Tables 17a, 17b).

Insects: jack pine budworm, white pine weevil, eastern pine shoot borer, jack pine tip beetle, jack pine sawflies and Swaine jack pine sawfly.

Diseases: Armillaria root rot, stem rusts, western gall rust, pine needle rust, tar spot needle cast.

Insects not found in the current survey included jack pine budworm and jack pine sawflies.

Moderate-to-severe damage by white pine weevil was noted in four plantations in the survey. Low levels of eastern pine shoot borer, jack pine tip beetle, and Swaine jack pine sawfly were also observed at various locations.

Light-to-moderate levels of the foliar diseases pine needle rust and tar spot needle cast were observed in approximately two-thirds of the plantations. Western gall rust, which has become a problem in the region, was found in six areas. The most damage from gall rust was found in Ben Nevis Twp, Kirkland Lake District, where 32.6% of the trees were severely affected. Trace levels of Armillaria root rot were found in three plantations.

Table 17a. Summary of the results of a special survey of 18 jack pine plantations or high-value stands in the region in 1988 (counts based on the examination of 150 or 300 randomly selected trees at each location).

Location (Twp)	Esti- mated stand area (ha)	Esti- mated no. of trees per ha	Height class (m)	Jack pine budworm		White pine weevil	Eastern pine shoot borer		Jack pine tip beetle	Swaine jack pine sawfly	
				Trees affected (%)	Defoliation (%)	Trees affected (%)	Leaders attacked (%)	Laterals attacked (%)	Leaders attacked (%)	Trees affected (%)	Defoliation (%)
<u>Chapleau District</u>											
Nimitz	134	2,500	<2 m	0	0	14.7	2	2	0	0	0
Edith	40	1,500	>6 m	0	0	0	0	0	0	0	0
Bliss <sup>a</sup>	75	2,500	2.0-6.0 m	0	0	0.3	3.0	0	0	0	0
Gilliland <sup>a</sup>	20	2,500	2.0-6.0 m	0	0	1.3	4.3	0	0	0	0
<u>Gogama District</u>											
Invergarry	10	2,500	<2 m	0	0	9.3	1.3	0	0	0	0
Vrooman	10	2,500	>6 m	0	0	0	0.6	1.2	0	0	0
<u>Kirkland Lake District</u>											
Cane	10	2,500	>6 m	0	0	1.3	0	0	0.7	0.7	2
Ingram	10	2,500	>6 m	0	0	0.7	0	0	2.7	0	0
Arnold	10	2,500	<2 m	0	0	6.7	0.7	2.0	0.7	0	0
Ben Nevis	10	2,500	<2 m	0	0	0.7	0.7	0	0	0	0
Gross <sup>a</sup>	15	2,500	2.0-6.0 m	0	0	14.0	3.3	4.0	4.0	0	0
<u>Timmins District</u>											
Robb <sup>a</sup>	50	2,500	2.0-6.0 m	0	0	0.7	3.3	0	0	0	0
<u>Cochrane District</u>											
Freele	3	1,600	<2 m	0	0	0.7	0	0	0	0	0
Sheldon <sup>a</sup>	93	1,600	2.0-6.0 m	0	0	0	0	0	0	0	0
Dempsey	10	1,600	>6 m	0	0	0	0	0	0	0	0
<u>Hearst District</u>											
Cross <sup>a</sup>	220	2,500	2.0-6.0 m	0	0	0	0	0	0	0	0
Larkin	5	2,500	<2 m	0	0	0	0	0	0	0	0
<u>Kapuskasing District</u>											
Fauquier	3	1,500	>6 m	0	0	0	0	0	0	0	0

<sup>a</sup> 300-tree sample

Table 17b. Summary of the results of a special survey of 18 jack pine plantations or high-value stands in the region in 1988 (counts based on the examination of 150 or 300 randomly selected trees at each location).

Location (Twp)	Esti- mated stand area (ha)	Esti- mated no. of trees per ha	Height class (m)	Armillaria	Stem rusts		Western gall rust		Pine	Tar spot
				root rot	Trees	Trees severely	Trees	Trees severely	needle rust	needle cast
				Trees affected (%)	Trees affected (%)	Trees severely affected (%)	Trees affected (%)	Trees severely affected (%)	Avg foliar damage (%)	Avg foliar damage (%)
<u>Chapleau District</u>										
Nimitz	134	2,500	<2 m	0	0	0	8	5.3	6.5	15
Edith	40	1,500	>6 m	0	0	0	14	1.3	0	2.7
Bliss <sup>a</sup>	75	2,500	2.0-6.0 m	0	0	0	0	0	8.7	9.3
Gilliland <sup>a</sup>	20	2,500	2.0-6.0 m	0	0	0	0	0	0	0
<u>Gogama District</u>										
Invergarry	10	2,500	<2 m	0.7	0	0	8.9	7.6	2.0	10.5
Vrooman	10	2,500	>6 m	0	0	0	24.0	0	6.5	6.5
<u>Kirkland Lake District</u>										
Cane	10	2,500	>6 m	0	0	0	0	0	0	12.0
Ingram	10	2,500	>6 m	0	0	0	0	0	2.0	5.0
Arnold	10	2,500	<2 m	0.7	0	0	0	0	1.0	0
Ben Nevis	10	2,500	<2 m	0	0	1.3	32.6	32.6	0	0
Gross <sup>a</sup>	10	2,500	2.0-6.0 m	0	0	0	0	0	0	5.0
<u>Timmins District</u>										
Robb <sup>a</sup>	50	2,500	2.0-6.0 m	0	0	0	0	0	0	0
<u>Cochrane District</u>										
Freele	3	1,600	<2.0 m	0.7	0	0	4.0	2.7	0	16.0
Sheldon <sup>a</sup>	93	1,600	2.0-6.0 m	0	0	0	0	0	0	0
Dempsey	10	1,600	>6.0 m	0	0	0	0	0	0	0
<u>Hearst District</u>										
Cross <sup>a</sup>	220	2,500	2.0-6.0 m	0	0	0	0	0	5	0.3
Larkin	5	2,500	<2 m	0	0	0	0	0	5	0
<u>Kapuskasing District</u>										
Fauquier	3	1,500	>6 m	0	0	0	0	0	0	0

<sup>a</sup> 300-tree sample

## Surveys of Nurseries

There are three provincial tree nurseries located in the region: one each in Chapleau, Gogama and Kirkland Lake districts. Contacts were maintained with nursery personnel, and surveys were conducted on a weekly or biweekly basis. Rangers were available if a problem arose that needed immediate attention.

At the Chapleau Nursery fewer than 1% of the black spruce planted in the spring of 1987 were affected with *Cylindrocladium floridanum* Sob. & C.P. Seym. This disease caused some mortality but, because of its low levels, it was not considered a problem. One jack pine seedling had its leader attacked by jack pine tip beetle, but again, because damage was minimal, it was not a problem.

Mature jack pine just outside the Gogama nursery are heavily infected with western gall rust. This gall rust may become a problem as it is possible for it to be transferred to jack pine seedlings. The infection will not become evident until the trees have been transplanted, as the fruiting does not appear until three or four years after infection. In addition, 1-2% of young white spruce in a seed orchard have been killed by armillaria root rot in the past couple of years.

At Swastika Nursery most damage was found on windbreak trees and in plantations. In jack pine plantations, white pine weevil and jack pine tip beetle were commonly found. Also found at low levels were tar spot needle cast and jack pine resin midge (*Cecidomyia resinicola* [O.S.]). Low levels of white pine blister rust were found scattered in the nursery. Trembling aspen trees growing in the nursery were infected with ink spot of aspen and shoot blight. As was the case throughout the region, severe defoliation of aspen and willow was caused by gray willow leaf beetle. Finally, two other pests, *Nycteola cinereana* N. & D. and *Lepyrus alternans* Casey, were found at low levels under container stock.

## Gypsy Moth Pheromone Traps

Pheromone traps were set out in the 12 provincial parks and one private park in the region to detect the presence of gypsy moth (*Lymantria dispar* [L.]). Ten traps were hung in each park in which positive catches had been made in 1987, viz., Fushimi Provincial Park, Hearst District, and Kettle Lakes Provincial Park, Timmins District. Two traps were deployed in each of the remaining parks. In 1988, male adult moths were caught in Esker Lakes Provincial Park and Kap-kig-i-wan Provincial Park, Kirkland Lake District. All other results were negative.

### Acid Rain National Early Warning System

As in previous years, the three study plots established in 1984 and 1985 were resampled in 1988. All the trees on the 400 m<sup>2</sup> plots were examined and any damage was recorded. Ground-cover plants were examined and compared with those collected in previous years. No damage was observed in any plot except for low levels of tar spot needle cast and western gall rust in the plot in Deans Township, Chapleau District.

Aerial photos are taken annually to detect any changes to the stand.

### Jack Pine Seed and Cone Pests

Two areas were sampled to assess the types of insects and diseases present and their impact on cone and seed crops (Table 18). One hundred 2-year-old cones from a minimum of three trees representing the total length of the cone-bearing crown were collected during the first week of July. The sample was forwarded to the Great Lakes Forestry Centre for dissection and analysis. Damage found in 1988 was generally low and insignificant.

Table 18. Summary of jack pine seed and cone damage at two locations in the region in 1988.

Location (Twp)	No. of cones examined	Damaged cones (%)	Seed loss within damaged cones (%)	Principal cause of seed loss (in order of importance)
<u>Chapleau District</u>				
Hong Kong	100	6	1.4	Lepidoptera Resseliella sp.
<u>Hearst District</u>				
Cross	100	7	2.9	Unknown Resseliella sp.

# Pinewood Nematode, *Bursaphelenchus xylophilus* (Steiner & Bührer) Nickle

As a result of sampling conducted in recent years it has been determined that this pest is present throughout the region on dead and dying jack pine. Therefore, work in 1988 was concentrated in other areas. Balsam fir (*Abies balsamea* [L.] Mill.) and spruce were sampled to determine if the "r" or "m" form of this pest was present (Fig. 6). Also, trees in known infection centers trees were marked as either dead or alive. The purpose of these designations was to determine if damage is spreading and, if it is, to document the rate of spread.

Work is also under way to determine possible vectors of the nematode. Fourteen samples of sawyer beetles and 21 samples of bark beetles were collected to determine if they were carrying the nematode (Fig. 7). These samples were submitted to experts at Memorial University in Newfoundland for analysis, but the results are not yet available.

## Climatic Data

Weather records for three stations, those at the Kapuskasing, Earlton and Chapleau airports, have been included in this report (Table 19). Weather plays an important part in insect development and the spread of infectious diseases, and is the cause of many non-infectious conditions such as frost and hail damage, winter drying and drought.

Table 19. Summary of mean temperature and total precipitation at three locations in the region in 1988.

Month	Mean temp. 1988 (°C)	Normal temp. (°C)	Deviation from norm temp. (°C)	Total precip. (mm)	Normal precip. (mm)	Deviation from norm precip. (mm)
<u>Earlton</u>						
Jan.	-15.2	-16.3	+1.1	60.7	56.4	+4.3
Feb.	-16.6	-14.1	-2.5	77.1	47.2	+29.9
Mar.	-8.0	-7.6	-0.4	50.5	58.0	-7.5
Apr.	2.3	1.9	+0.4	110.5	50.0	+60.5
May	11.8	9.8	+2.0	63.4	61.3	+2.1
June	14.6	15.2	-0.6	60.3	89.2	-28.9
July	19.8	17.7	+2.1	34.6	80.8	-46.2
Aug.	17.6	16.2	+1.4	165.5	83.4	+82.1
Sept.	11.5	11.1	+0.4	66.3	99.1	-32.8
Oct.	3.6	5.4	-1.8	65.4	70.0	-4.6
Nov.	-0.4	-2.5	+2.1	84.6	70.6	+14.0
Dec.	-13.7	-12.6	-1.1	49.9	65.3	-15.4

(cont'd)

Table 19. Summary of mean temperature and total precipitation at three locations in the region in 1988 (concl.).

Month	Mean temp. 1988 (°C)	Normal temp. (°C)	Deviation from norm temp. (°C)	Total precip. (mm)	Normal precip. (mm)	Deviation from norm precip. (mm)
<u>Chapleau</u>						
Jan.	-16.1	-16.9	+0.8	76.2	46.9	+29.3
Feb.	-16.3	-15.8	-0.5	43.8	34.5	+9.3
Mar.	-8.3	-8.6	+0.3	68.6	56.2	+12.4
Apr.	2.5	0.6	+1.9	4.8	59.3	-51.5
May	11.1	8.6	+2.5	57.4	73.8	-16.4
June	14.8	14.3	+0.2	57.8	100.4	-42.6
July	19.1	16.8	+2.3	41.6	81.8	-40.2
Aug.	16.8	15.4	+1.4	170.0	86.2	+83.8
Sept.	11.0	10.4	+0.6	79.6	101.5	+21.9
Oct.	*	4.9		*	75.7	
Nov.	*	-3.5		*	64.2	
Dec.	*	-12.8		*	53.5	
<u>Kapuskasing</u>						
Jan.	-18.7	-18.6	-0.1	57.7	53.6	+4.1
Feb.	-19.1	-16.2	-2.9	55.2	43.0	+12.2
Mar.	-10.3	-9.4	-0.9	74.2	55.4	+18.8
Apr.	1.4	0.5	+0.9	42.8	53.2	-10.4
May	10.2	8.3	+1.9	69.8	74.3	-4.5
June	13.0	14.1	-1.1	82.0	84.7	-2.7
July	19.0	16.8	+2.2	57.6	96.3	-38.7
Aug.	15.7	15.3	+0.4	179.2	92.5	+86.7
Sept.	10.6	10.0	+0.6	85.5	94.4	-8.9
Oct.	2.3	4.4	-2.1	60.9	77.4	-16.5
Nov.	-2.4	-4.4	+2.0	127.1	80.1	+47.0
Dec.	-16.4	-14.7	-1.7	61.6	53.3	+8.3

\* Data not available at the time of publication

## NORTHERN REGION




Figure 6. Pinewood nematode (*Bursaphelenchus xylophilus* [Steiner and Bührer] survey. Areas sampled in 1988:

jack pine ▲  
balsam fir ○



## NORTHERN REGION




Figure 7. Pinewood nematode (*Bursaphelenchus xylophilus* [Steiner and Bührer] Nickle) vector survey, 1988.

Locations sampled: *Monochamus* spp. ▲

bark beetles ■